arXiv:1211.1931v1 [math.AG] 8 Nov 2012

Modular Subgroups, Dessins d'Enfants and Elliptic K3 Surfaces

Yang-Hui He[1], John McKay[2] & James Read[3]

[1]*Department of Mathematics, City University, London,*
Northampton Square, London EC1V 0HB, UK;
School of Physics, NanKai University, Tianjin, 300071, P.R. China;
Merton College, University of Oxford, OX1 4JD, UK
yang-hui.he@merton.ox.ac.uk

[2]*Department of Mathematics and Statistics,*
Concordia University, 1455 de Maisonneuve Blvd. West,
Montreal, Quebec, H3G 1M8, Canada
mckay@encs.concordia.ca

[3]*Oriel College, University of Oxford, OX1 4EW, UK*
james.read@oriel.ox.ac.uk

Abstract

We consider the 33 genus zero, torsion-free modular subgroups, computing ramification data and Grothendieck's *dessins d'enfants*. In the particular case of the index 36 subgroups, the corresponding Calabi-Yau threefolds are identified, in analogy with the index 24 cases being associated with K3 surfaces. In a parallel vein, we study the 112 semi-stable elliptic fibrations over $\mathbb{P}^1$ as extremal K3 surfaces with six singular fibres. In each case, the corresponding modular subgroup is identified by showing its generators.

Contents

1 Introduction 3

2 Torsion-Free, Genus Zero Congruence Subgroups 4

2.1 The Modular Group and Congruence Subgroups 4

2.1.1 The 33 Subgroups 6

2.2 Modular Surfaces, Ramification Data and j-Functions 8

2.3 Dessins d'Enfants: j-Functions as Belyi Maps 9

2.3.1 Dessins for the 33 Subgroups 10

2.4 Index 36 Subgroups and Calabi-Yau Threefolds 14

2.4.1 Example: $\Gamma_0(2) \cap \Gamma(3)$ 16

3 Semi-Stable Elliptic K3 Surfaces 17

3.1 Permutation Representations and Cartographic Groups 18

3.1.1 Cartographic Groups for the 112 K3 Surfaces 19

3.2 Modular Subgroup Generators from Cartographic Groups 26

3.3 Subgroup Presentations: Free Groups on Five Generators 28

3.4 Generators for the Modular Subgroups 31

3.4.1 Catalogue of Generators as Explicit 2×2 Matrices 32

4 Conclusions and Outlook 41

A Dessins from the Miranda-Persson Table 41

1 Introduction

The congruence subgroups of the modular group $\Gamma = PSL(2;\mathbb{Z})$ play a key role in many branches of modern mathematics. Whereas the upper half of the complex plane $\mathcal{H}$, when adjoining cusps on the real line, can quotient Γ to give a Riemann sphere, it is an interesting question as for which subgroups a similar sphere arises. Such a subgroup is called genus zero. Inspired by Moonshine, the authors of [1] looked into the classification problem of congruence, genus zero, torsion free subgroups of the modular group and found these to be rather rare: there are only 33 of them, of finite index μ equaling 6,12,24,36,48, or 60.

Therefrom emerges many remarkable connections to geometry, graph theory, number theory and physics [2, 3, 4]. First, we naturally extend the modular group action to a so-called modular surface for each of the groups whose Euler number is precisely μ. These turn out to be elliptically fibred over the genus zero $\mathbb{P}^1$ formed by quotienting $\mathcal{H}$ by the group and are, in fact, semi-stable in the sense that all singular fibres are of Kodaira type I_n. Moreover, there are exactly $h = \frac{\mu}{6} + 2$ such singular fibres, and the tuple $\{n_1, \ldots, n_h\}$ is called the cusp widths.

Next, one can construct the Schreier coset graphs for these groups because they have finite index. They turn out to be finite trivalent graphs of non-trivial topology. We can interpret them as clean cubic dessin d'enfants in the sense of Groethendieck. Focus has been made on the index $\mu = 24$ cases because they are K3 surfaces [3] and on the $\mu = 12$ cases because they have interesting associated modular forms [2]. One can check that the Belyi rational maps for the dessins are precisely the elliptic j-invariants of the Weierstraß models for the modular surfaces with base $\mathbb{P}^1$ and we will complete this story in the present paper.

In a slightly different but obviously related direction, the classification of extremal, semi-elliptic K3 surfaces with 6 singular fibres was wonderfully performed by Miranda and Persson in [10], and the associated dessins nicely compiled in [11]. It is therefore a natural question whether they can be affiliated with subgroups of the modular group, not necessarily congruence. Looking at these dessins, one see them to be trivalent and clean, and since these can be interpreted as the Schreier coset graphs of a subgroup of the modular group, the answer must be affirmative. We shall undertake this task, using the dessins, to find the generators for all of them.

The organization of the paper is as follows. In §2, we introduce the 33 congruence, torsion-free, genus zero subgroups, especially their j-invariants, dessins and Schreier coset graphs. We will see that as the index 24 cases are attached to K3 surfaces, so too can the index 36 cases be associated to Calabi-Yau threefolds. Then, in §3, we turn to the 112 extremal semi-stable K3 surfaces, studying their dessins and interpreting them as coset graphs of some subgroup. Via a discussion of the elliptic j-invariants as Belyi maps and the so-called cartographic groups, we find the generators for these subgroups. Finally, we conclude with some outlook and prospects.

2 Torsion-Free, Genus Zero Congruence Subgroups

In this section, we first recall some standard notation regarding the modular group Γ and its congruence subgroups. We focus specifically on the case of the *torsion-free, genus zero, congruence subgroups*; there are 33 such groups as classified by [1]. For each group, we will compute the ramification data, and draw the dessin d'enfant which encodes this data. The case of index 24 and 36 are particularly interesting: the former can be associated to K3 surfaces and we will demonstrate, using the j-invariants, that each of the index 36 subgroups can be identified with a specific Calabi-Yau threefold.

2.1 The Modular Group and Congruence Subgroups

First, the central object of our study is the *modular group*

$$\Gamma \equiv \Gamma(1) = \mathrm{PSL}(2;\mathbb{Z}) = \mathrm{SL}(2;\mathbb{Z})/\left\{\pm I\right\} \ . \tag{2.1}$$

This is the group of linear fractional transformations $z \to \frac{az+b}{cz+d}$, with $a,b,c,d \in \mathbb{Z}$ and $ad-bc=1$. It is generated by the transformations T and S defined by:

$$T\tau = \tau + 1 \quad , \quad S\tau = -1/\tau \ . \tag{2.2}$$

One presentation of Γ is $\left\langle S,T|S^2 = (ST)^3 = I\right\rangle$, and we will later discuss the presentations of certain modular subgroups. 2×2 matrices for S and T are as follows:

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad , \quad S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \ . \tag{2.3}$$

Letting $x = S$ and $y = ST$ denote the elements of order 2 and 3 respectively, we see that Γ is the free product of the cyclic groups $C_2 = \langle x|x^2 = I\rangle$ and $C_3 = \langle y|y^3 = I\rangle$. It follows that 2×2 matrices for x and y are:

$$x = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad , \quad y = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix} \ . \tag{2.4}$$

Cayley Graphs: Given the free product structure of Γ, we see that its Cayley graph is an infinite free trivalent tree, but with each node replaced by an oriented triangle. Now, for any *finite index* subgroup H of Γ, we can quotient the Cayley graph to arrive at a finite cubic graph by associating nodes to right cosets and edges between cosets which are related by action of a group element. In other words, this graph encodes the permutation representation of Γ acting on the right cosets of H. This is called a **Schreier coset graph** and will occupy a central role in our ensuing discussions. It is a remarkable fact that *all* trivalent graphs arise as the coset graph of some finite index subgroup of Γ.

Congruence Subgroups: The most important subgroups of Γ are the so-called congruence subgroups, defined by having the entries in the generating matrices S and T obeying some modular arithmetic. Some congruence subgroups of particular note are the following:

- Principal congruence subgroups:

$$\Gamma\left(m\right) := \{A \in \mathrm{SL}(2;\mathbb{Z})\ ,\ A \equiv \pm I \bmod m\} \,/\, \{\pm I\}\,;$$

- Congruence subgroups of level m: subgroups of Γ containing $\Gamma\left(m\right)$ but not any $\Gamma\left(n\right)$ for $n < m$;
- Unipotent matrices:

$$\Gamma_1\left(m\right) := \left\{A \in \mathrm{SL}(2;\mathbb{Z})\ ,\ A \equiv \pm \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \bmod m\right\} / \{\pm I\}\,;$$

- Upper triangular matrices:

$$\Gamma_0\left(m\right) := \left\{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma\ ,\ c \equiv 0 \bmod m\right\} / \{\pm I\}\,.$$

In [3], our attention was drawn to a particular family of subgroups of Γ: the so-called *genus zero, torsion-free* congruence subgroups. To recapitulate, by *torsion-free* we mean that the subgroup contains no element of finite order other than the identity. To explain *genus zero*, first recall that the modular group acts on the upper half-plane $\mathcal{H} := \{\tau \in \mathbb{C}\ ,\ \mathrm{Im}\left(\tau\right) > 0\}$ by linear fractional transformations $z \to \frac{az+b}{cz+d}$. $\mathcal{H}$ gives rise to a compactification $\mathcal{H}^*$ when adjoining *cusps*, which are points on $\mathbb{R}\cup\infty$ *fixed* under some parabolic element (i.e. an element in Γ not equal to the identity which nevertheless has trace 2).

The quotient $\mathcal{H}^*/\Gamma$ is a compact Riemann surface of genus 0, i.e., a sphere. It turns out that with the addition of appropriate cusp points, the extended upper half plane $\mathcal{H}^*$ factored by various congruence subgroups will also be compact Riemann surfaces, possibly of higher genus. Such a Riemann surface, as a complex algebraic variety, is called a **modular curve**. The genus of a subgroup of the modular group is the genus of the modular curve produced in this way.

Using the Riemann-Hurwitz theorem, the genus of the modular curve can be shown to be:

$$g_C = 1 + \frac{\mu}{12} - \frac{\nu_2}{4} - \frac{\nu_3}{3} - \frac{\nu_\infty}{2} \ , \tag{2.5}$$

where μ is the index of the subgroup, ν_2, ν_3 are the number of inequivalent elliptic points of order 2 and 3 respectively, and ν_∞ is the number of inequivalent cusp points.

Our focus will be on genus zero, torsion-free subgroups, thus we have that

$$g_C = \nu_2 = \nu_3 = 0 \ , \tag{2.6}$$

whence, by (2.5), we have a further constraint that $\mu = 6\nu_\infty - 12$, so that the index of all our subgroups of concern is a multiple of 6. This means that the index of the subgroup in question also constrains ν_∞, so that for a subgroup of index $6n$, we have $\nu_\infty = n + 2$.

2.1.1 The 33 Subgroups

As classified in [1], the genus zero torsion-free congruence subgroups of the modular group are very rare. There are only 33 of them:

- $\Gamma(n)$ with $n = 2, 3, 4, 5$;
- $\Gamma_0(n)$ with $n = 4, 6, 8, 9, 12, 16, 18$;
- $\Gamma_1(n)$ with $n = 5, 7, 8, 9, 10, 12$;
- The intersections $\Gamma_0(a) \cap \Gamma(b)$ for $\{a, b\} = \{4, 2\}, \{3, 2\}, \{8, 2\}, \{2, 3\}, \{25, 5\}$; $\Gamma_1(8) \cap \Gamma(2)$ and $\Gamma_0(16) \cap \Gamma_1(8)$;
- The congruence subgroups

  $$\Gamma\left(m; \frac{m}{d}, \epsilon, \chi\right) := \left\{ \pm \begin{pmatrix} 1 + \frac{m}{\epsilon\chi}\alpha & d\beta \\ \frac{m}{\chi}\gamma & 1 + \frac{m}{\epsilon\chi}\delta \end{pmatrix} \ , \ \gamma \equiv \alpha \bmod \chi \right\}$$

  with $(m, d, \epsilon, \chi) = (8, 2, 1, 2)$, $(12, 2, 1, 2)$, $(16, 1, 2, 2)$, $(27, 1, 3, 3)$, $(8, 4, 1, 2)$, $(9, 3, 1, 3)$, $(16, 2, 2, 2)$, $(24, 1, 2, 2)$, $(32, 1, 4, 2)$.

For reference, in Table 1, we organize these 33 subgroups according to the index and we will shortly explain the notation of the ramification data.

Index	Group	Ramification
6	$\Gamma(2)$	$\left\{\begin{matrix}3^2\\2^3\\2^3\end{matrix}\right\}$
6	$\Gamma_0(4)$	$\left\{\begin{matrix}3^2\\2^3\\1^2,4\end{matrix}\right\}$

Index	Group	Ramification
12	$\Gamma(3)$	$\left\{\begin{matrix}3^4\\2^6\\3^4\end{matrix}\right\}$
12	$\Gamma_0(4)\cap\Gamma(2)$	$\left\{\begin{matrix}3^4\\2^6\\2^2,4^2\end{matrix}\right\}$
12	$\Gamma_1(5)$	$\left\{\begin{matrix}3^4\\2^6\\1^2,5^2\end{matrix}\right\}$
12	$\Gamma_0(6)$	$\left\{\begin{matrix}3^4\\2^6\\1,2,3,6\end{matrix}\right\}$
12	$\Gamma_0(8)$	$\left\{\begin{matrix}3^4\\2^6\\1^2,2,8\end{matrix}\right\}$
12	$\Gamma_0(9)$	$\left\{\begin{matrix}3^4\\2^6\\1^3,9\end{matrix}\right\}$

Index	Group	Ramification
24	$\Gamma(4)$	$\left\{\begin{matrix}3^8\\2^{12}\\4^6\end{matrix}\right\}$
24	$\Gamma_0(3)\cap\Gamma(2)$	$\left\{\begin{matrix}3^8\\2^{12}\\2^3,6^3\end{matrix}\right\}$
24	$\Gamma_1(7)$	$\left\{\begin{matrix}3^8\\2^{12}\\1^3,7^3\end{matrix}\right\}$
24	$\Gamma_1(8)$	$\left\{\begin{matrix}3^8\\2^{12}\\1^2,2,4,8^2\end{matrix}\right\}$
24	$\Gamma_0(8)\cap\Gamma(2)$	$\left\{\begin{matrix}3^8\\2^{12}\\2^4,8^2\end{matrix}\right\}$
24	$\Gamma(8;4,1,2)$	$\left\{\begin{matrix}3^8\\2^{12}\\2^2,4^3,8\end{matrix}\right\}$
24	$\Gamma_0(12)$	$\left\{\begin{matrix}3^8\\2^{12}\\1^2,3^2,4,12\end{matrix}\right\}$
24	$\Gamma_0(16)$	$\left\{\begin{matrix}3^8\\2^{12}\\1^4,4,16\end{matrix}\right\}$
24	$\Gamma(16;16,2,2)$	$\left\{\begin{matrix}3^8\\2^{12}\\1^2,2^3,16\end{matrix}\right\}$

Index	Group	Ramification
36	$\Gamma_0(2)\cap\Gamma(3)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\3^4,6^4\end{matrix}\right\}$
36	$\Gamma_1(9)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\1^3,3^2,9^3\end{matrix}\right\}$
36	$\Gamma(9;3,1,3)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\3^6,9^2\end{matrix}\right\}$
36	$\Gamma_1(10)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\1^2,2^2,5^2,10^2\end{matrix}\right\}$
36	$\Gamma_0(18)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\1^3,2^3,9,18\end{matrix}\right\}$
36	$\Gamma(27;27,3,3)$	$\left\{\begin{matrix}3^{12}\\2^{18}\\1^6,3,27\end{matrix}\right\}$

Index	Group	Ramification
48	$\Gamma_1(8)\cap\Gamma(2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\2^4,4^2,8^4\end{matrix}\right\}$
48	$\Gamma(8;2,1,2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\4^8,8^2\end{matrix}\right\}$
48	$\Gamma_1(12)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\1^2,2,3^2,4^2,6,12^2\end{matrix}\right\}$
48	$\Gamma(12;6,1,2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\2^4,4,6^4,12\end{matrix}\right\}$
48	$\Gamma_0(16)\cap\Gamma_1(8)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\1^4,2^2,4^2,16^2\end{matrix}\right\}$
48	$\Gamma(16;8,2,2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\2^8,16^2\end{matrix}\right\}$
48	$\Gamma(24;24,2,2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\1^4,3^4,8,24\end{matrix}\right\}$
48	$\Gamma(32;32,4,2)$	$\left\{\begin{matrix}3^{16}\\2^{24}\\1^8,8,32\end{matrix}\right\}$

Index	Group	Ramification
60	$\Gamma(5)$	$\left\{\begin{matrix}3^{20}\\2^{30}\\5^{12}\end{matrix}\right\}$
60	$\Gamma_0(25)\cap\Gamma_1(5)$	$\left\{\begin{matrix}3^{20}\\2^{30}\\1^{10},25^2\end{matrix}\right\}$

Table 1: *The 33 genus zero, torsion free, congruence subgroups of the modular group, organized by index (which can only be 6, 12, 24, 36, 48 or 60). The ramification data for each group is defined in* (2.9).

As a point of interest, we note here that $\Gamma(2)$ and $\Gamma_0(4)$ are conjugate subgroups. By this, we mean that there exists a matrix M such that, for all elements $A_{\Gamma(2)}$ of $\Gamma(2)$ there is an element $B_{\Gamma_0(4)}$ of $\Gamma_0(4)$ satisfying

$$M^{-1}A_{\Gamma(2)}M = B_{\Gamma_0(4)} \tag{2.7}$$

The matrix M can be found by choosing a generator of $\Gamma(2)$ and a generator of $\Gamma_0(4)$, substituting into the above equation, and solving. For example, using the procedure in §3.4, we find that a set of generators for $\Gamma(2)$ is $\left\{\begin{pmatrix}-1 & 0\\ 2 & -1\end{pmatrix}, \begin{pmatrix}-1 & 2\\ 0 & -1\end{pmatrix}\right\}$. Applying conjugation by the matrix $M = \begin{pmatrix}0 & 1\\ 2 & 1\end{pmatrix}$, we obtain the matrices $\left\{\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}, \begin{pmatrix}-3 & -1\\ 4 & 1\end{pmatrix}\right\}$. These can again be easily checked to be the generators of $\Gamma_0(4)$.

2.2 Modular Surfaces, Ramification Data and j-Functions

As detailed in [3], to each torsion-free, genus zero modular subgroup corresponds a distinct **Shioda modular surface** [5]. We briefly recall how this is constructed [4]. First, we extend the action of any subgroup G of Γ on $\mathcal{H}$ to an action

$$\mathcal{H} \times \mathbb{C} \ni (\tau, z) \longrightarrow \left(\gamma\tau, \frac{z + m\tau + n}{c\tau + d}\right) , \tag{2.8}$$

for $\gamma = \begin{pmatrix}a & b\\ c & d\end{pmatrix} \in G$ and $(m, n) \in \mathbb{Z}^2$. Next, the quotient of $\mathcal{H} \times \mathbb{C}$ by the above automorphism defines a surface equipped with a morphism to the modular curve C arising from the quotient of $\mathcal{H}$ by $\tau \to \gamma\tau$. The fiber over the image of this morphism to the modular curve is generically an elliptic curve corresponding to the lattice $\mathbb{Z} \oplus \mathbb{Z}\tau$. What we have therefore is a complex surface which is an elliptic fibration over the modular curve associated to G. This is the modular surface.

In our present case, because the base modular curves are genus zero (being the Riemann sphere $\mathbb{P}^1_C$), our modular surfaces will be elliptic fibrations over $\mathbb{P}^1_C$. It turns out the Euler number of this surface is the index μ of the group H. Hence, the astute reader will recognize that the index 24 groups should give us K3 surfaces, in particular elliptic K3 surfaces (we remark that $\Gamma_1(7)$, in particular, was realized by Tate to be related to a K3 surface). We will return to this point later, especially to the observation that index 36 groups, by analogy, give Calabi-Yau threefolds.

Furthermore, the elliptic modular surface is *semi-stable* in the sense that all the singular fibres are of Kodaira type I_n; the set of integers $(n_1, n_2, \ldots, n_k)$ denoting the indices of the singular fibres are the **cusp widths**. For simplicity, we arrange this set in increasing order and group repeated entries by the short-hand $n_i^{a_i}$ with a_i being the multiplicity. Now, we are ready to introduce the concept of **ramification data** associated with a subgroup (and its modular surface), it is a triplet of integer vectors

$$\left\{ \begin{array}{c} 3^V \\ 2^E \\ n_1^{a_1}, n_2^{a_2}, \ldots, n_k^{a_k} \end{array} \right\}, \qquad \begin{array}{rcl} V & = & \text{number of nodes in the Schreier coset graph} \\ E & = & \text{number of edges} \\ \{n_i^{a_i}\} & = & \text{cusp widths of the elliptic modular surface} \end{array} \tag{2.9}$$

Note that the 3 reminds us that all nodes are trivalent and 2, that each edge links two nodes. We have marked this ramification data in Table 1.

Indeed, given an elliptic fibration, we can write down a Weierstraß equation, from which we can then extract a (Klein) j-function. Because of the fibration over $\mathbb{P}^1_C$, the j-function will explicitly depend on the projective coordinate of this base Riemann sphere and can thus be treated as a rational map from the base $\mathbb{P}^1_C$ to its range $\mathbb{P}^1$. All of these j-functions are given in [1] and certain simplifications, in [3, 4]. Generalizing [3, 10], we find the nice fact that

PROPOSITION 1 *The j-function $j\left(t \in \mathbb{P}^1_C\right) \mapsto \mathbb{P}^1$ is ramified at only $(0, 1, \infty)$.*

In other words, the only points $t \in \mathbb{P}^1_C$ such that $\frac{\mathrm{d}}{\mathrm{d}t} j\left(t\right)|_{\tilde{t}} = 0$ are such that $j(\tilde{t}) = 0, 1$ or ∞. Moreover, the order of vanishing of the Taylor series at t is the ramification index (we remark that depending on convention, we may need to appropriately normalize to the j-function by a factor of 1728).

Specifically, for our 33 subgroups, we find that for a group of index $6n$, there are $2n$ pre-images of 0 and the ramification indices of all these marked points is 3; there are $3n$ pre-images of 1 and ramification indices of all these marked points is 2; there are $n+2$ pre-images of ∞, the ramification indices of which are given as the cusp widths in Table 1.

2.3 Dessins d'Enfants: j-Functions as Belyi Maps

Maps to $\mathbb{P}^1$ ramified only at $(0, 1, \infty)$ are called **Belyi maps** and can be represented graphically as Grothendieck's *dessins d'enfants*. To draw such a dessin is simple: given ramification data $\left\{ \begin{array}{c} \{r_0\left(i\right)\} \\ \{r_1\left(i\right)\} \\ \{r_\infty\left(i\right)\} \end{array} \right\}$ specifying the ramification indices at the various pre-images of 0, 1 and infinity, one marks one white node for the i-th pre-image of 0, with $r_0\left(i\right)$ edges emanating therefrom; similarly, one marks one black node for the j-th pre-image of 1, with $r_1(j)$ edges. Thus we have a **bipartite graph** embedded on a Riemann sphere, with W white nodes and B black nodes. Now we connect the nodes with the edges, joining only black with white, such that we have I faces, each being a polygon with $2r_\infty(k)$ sides. Dessins for all the 33 torsion-free, genus zero modular subgroups are presented in §2.3.1.

The Schreier coset graphs for the 33 subgroups are discussed and drawn explicitly in [1]. Interestingly, there is a direct connection between these and the dessins d'enfants. In our case, let Γ^* be the torsion-free, genus zero modular subgroup in question. The coset graphs display the permutation action of generators x and y with the relations $x^2 = y^3 = I$ on each coset $\Gamma^* x_i$ of $PSL(2,\mathbb{Z}) = \bigcup_{i=1}^{\mu} \Gamma^* x_i$ where μ is the index of Γ^* in $PSL(2,\mathbb{Z})$.

These graphs have μ nodes, where μ is the index and takes the values $6, 12, 24, 36, 48, 60$. The graphs are built from simple edges (x) and triangles (y) which are assumed to be positively oriented [*].

The connection between the dessins d'enfants and the Schreier coset graphs can be stated as follows:

PROPOSITION 2 *The dessins can be constructed from the Schreier coset graphs by replacing each positively oriented triangle with a white node, and inserting a black node into every edge. Equivalently, the Schreier coset graphs can be constructed from the dessins by replacing each white node with a positively oriented triangle, and removing the black node from every edge.*

The process of inserting into an edge a bivalent node of a different colouring is standard in the study of dessins d'enfants; this gives rise to so-called *clean* dessins. The contraction of triangles into a node is a less obvious one and should correspond to taking a cube root of the polynomial factors in the Belyi map.

2.3.1 Dessins for the 33 Subgroups

In this section, we draw the dessins d'enfants for all 33 genus zero, torsion-free congruence subgroups using the ramification data in Table 1. Each dessin is labelled by its corresponding modular subgroup, and the unique cusp widths for that subgroup. The reader can visually confirm the aforementioned correspondence between these dessins and the Schreier coset graphs constructed in [1].

---

[*] If, instead, we were to choose all negatively oriented triangles, we would still obtain correct results encoding the same information. However, the matrix generators which we will present later in §3.4 would be transposed, and therefore not in line with the conventions for the modular subgroups of interest laid out in §2.1. Moreover, if we instead choose random triangle orientations, the relation of the generators in §3.4 to the forms of the matrices for the modular subgroups of interest given in §2.1 is non-trivial. This provides positive motivation for taking the triangles to be positively oriented.



$\Gamma(2)$
$2, 2, 2$



$\Gamma_0(4)$
$4, 1, 1$



$\Gamma(3)$
$3, 3, 3, 3$



$\Gamma_0(4) \cap \Gamma(2)$
$4, 4, 2, 2$



$\Gamma_1(5)$
$5, 5, 1, 1$



$\Gamma_0(6)$
$6, 3, 2, 1$



$\Gamma_0(8)$
$8, 2, 1, 1$



$\Gamma_0(9)$
$9, 1, 1, 1$



$\Gamma(4)$
$4, 4, 4, 4, 4, 4$



$\Gamma(8; 4, 1, 2)$
$8, 4, 4, 4, 2, 2$



$\Gamma_0(3) \cap \Gamma(2)$
$6, 6, 6, 2, 2, 2$



$\Gamma_0(12)$
$12, 4, 3, 3, 1, 1$



$\Gamma_1(8)$
$8, 8, 4, 2, 1, 1$



$\Gamma_0(8) \cap \Gamma(2)$
$8, 8, 2, 2, 2, 2$



$\Gamma_0\,(16)$
$16, 4, 1, 1, 1, 1$



$\Gamma\,(16; 16, 2, 2)$
$16, 2, 2, 2, 1, 1$



$\Gamma_1\,(7)$
$7, 7, 7, 1, 1, 1$



$\Gamma_0\,(2) \cap \Gamma\,(3)$
$6, 6, 6, 6, 3, 3, 3, 3$



$\Gamma_1\,(9)$
$9, 9, 9, 3, 3, 1, 1, 1$



$\Gamma\,(9; 3, 1, 3)$
$9, 9, 3, 3, 3, 3, 3, 3$



$\Gamma_1\,(10)$
$10, 10, 5, 5, 2, 2, 1, 1$



$\Gamma_0\,(18)$
$18, 9, 2, 2, 2, 1, 1, 1$



$\Gamma\,(27; 27, 3, 3)$
$27, 3, 1, 1, 1, 1, 1, 1$



$\Gamma_1(8) \cap \Gamma(2)$

8, 8, 8, 8, 4, 4, 2, 2, 2, 2



$\Gamma(8; 2, 1, 2)$

8, 8, 4, 4, 4, 4, 4, 4, 4, 4



$\Gamma_1(12)$

12, 12, 6, 4, 4, 3, 3, 2, 1, 1



$\Gamma(12; 6, 1, 2)$

12, 6, 6, 6, 6, 4, 2, 2, 2, 2



$\Gamma_0(16) \cap \Gamma_1(8)$

16, 16, 4, 4, 2, 2, 1, 1, 1, 1



$\Gamma(16; 8, 2, 2)$

16, 16, 2, 2, 2, 2, 2, 2, 2, 2



$\Gamma(24; 24, 2, 2)$

24, 8, 3, 3, 3, 3, 1, 1, 1, 1



$\Gamma(32; 32, 4, 2)$

32, 8, 1, 1, 1, 1, 1, 1, 1, 1



$\Gamma(5)$

$5,5,5,5,5,5,5,5,5,5,5,5$



$\Gamma_0(25) \cap \Gamma_1(5)$

$25,25,1,1,1,1,1,1,1,1,1,1$

2.4 Index 36 Subgroups and Calabi-Yau Threefolds

Since all the index 24 torsion-free, genus zero congruence subgroups have been associated in [1, 4, 11] to K3 surfaces - in particular elliptically fibred K3 surfaces over $\mathbb{P}^1$ - it is natural to enquire whether the index 36 groups can be associated with Calabi-Yau threefolds. We shall see that this is indeed the case, and the threefolds are a simple generalization of the K3 case, being now elliptic fibrations over $\mathbb{P}^2$.

First, let us recall some standard facts about elliptic Calabi-Yau threefolds. The classification of base surfaces B over which an elliptic fibration is a smooth Calabi-Yau threefold was performed in [16, 17]: they can be either Hirzebruch surfaces and their blowups; del Pezzo surfaces; or the Enriques surface. In particular, the local Weierstraß model looks like:

$$y^2 = x^3 + f(z_i)\,x + g(z_i) \subset \mathbb{P}^2_{[x,y]} \ , \tag{2.10}$$

with

$$f(z_i) \in \Gamma\left(\left(K_B^{-1}\right)^{\otimes 4}\right) \quad , \quad g(z_i) \in \Gamma\left(\left(K_B^{-1}\right)^{\otimes 6}\right) \ . \tag{2.11}$$

This is an elliptic curve in a $\mathbb{P}^2$ bundle over B, whose coordinates are z_i; the (x, y) are affine coordinates on the $\mathbb{P}^2$. The functions f and g are homogeneous polynomials in the base coordinates. Specifically, they should be considered as sections Γ of appropriate tensor powers of the anti-canonical bundle K_B^{-1} of B: for f, it is the fourth power and for g, the sixth. A nice summary discussion of this can be found, for example, in §3 of [16].

The j-invariant of a Weierstraß model is given by:

$$j\left(z_i\right) = 1728 \cdot \frac{4f\left(z_i\right)^3}{4f\left(z_i\right)^3 - 27g\left(z_i\right)^2} \ . \tag{2.12}$$

Here, we have marked explicitly the dependence on the base coordinates z_i. In particular, we can see that both the numerator and denominator are sections of $\left(K_B^{-1}\right)^{\otimes 12}$.

Let us return to the index 36 torsion-free, genus zero congruence subgroups. As mentioned, the j-invariants for these are computed and tabulated in [1]; these are rational functions in terms of the principal modulus t and we interpreted these as Belyi maps in our above discussions to construct the dessins. From j we can extract the corresponding polynomials $f\left(t\right)$ and $g\left(t\right)$ for each subgroup. We find that the $f\left(t\right)$ will be a degree 12 and $g\left(t\right)$, degree 18, polynomial in t. In general, for an index μ subgroup (recall again that $\mu = 6, 12, 24, 36, 48, 60$ for genus zero, torsion-free congruence subgroups), the degrees of f and g are $\mu/3$ and $\mu/2$.

Finally, let us recall a standard fact about $\mathbb{P}^2$ to complete the story. The anti-canonical bundle of the projective plane is $K_{\mathbb{P}^2}^{-1} = \mathcal{O}_{\mathbb{P}^2}\left(3\right)$, therefore sections of this would be degree 3 homogeneous polynomials in the coordinates. Therefore, if we choose our base B to be $\mathbb{P}^2$ with affine coordinates z_1, z_2, then $f\left(z_i\right)$ will be a degree $3 \times 4 = 12$ and $g\left(z_i\right)$ a degree $3 \times 6 = 18$ homogeneous polynomial in z_1, z_2. These are the right degrees to match those of $f\left(t\right)$ and $g\left(t\right)$ for the index 36 case! Therefore, setting a linear change of basis (normalizing the leading coefficient)

$$z_1 = t + a \quad , \quad z_2 = t + b \qquad a, b \in \mathbb{C} \tag{2.13}$$

and substituting into appropriately chosen $f\left(z_i\right)$ and $g\left(z_i\right)$ should give the required $f\left(t\right)$ and $g\left(t\right)$. Geometrically, this amounts to finding a hyper-plane $\mathbb{P}^1$ defined by (2.13) inside the base $B = \mathbb{P}^2$ on which our elliptically fibred Calabi-Yau threefold restricts to the required modular surface for index 36.

The analogous situation for the index 24 case is clear. There, we have K3 surfaces which are elliptic fibrations over a base $\mathbb{P}^1$: we know that the anti-canonical bundle is $K_{\mathbb{P}^2}^{-1} = \mathcal{O}_{\mathbb{P}^2}\left(2\right)$ whose sections are thus quadratics in the projective coordinate t. Hence, $f\left(t\right)$ and $g\left(t\right)$ should be of degree $2 \times 4 = 8$ and $2 \times 6 = 12$, which are indeed in accord with the aforementioned fact that the degrees should be respectively $\mu/3$ and $\mu/2$ with $\mu = 24$ here.

2.4.1 Example: $\Gamma_0(2) \cap \Gamma(3)$

Let us see how the above setup works for the specific groups in detail. The j-invariant for the group $\Gamma_0(2) \cap \Gamma(3)$ is [1]:

$$j(t) = \frac{\left((t^3+4)(t^3+6t^2+4)(t^6-6t^5+36t^4+8t^3-24t^2+16)\right)^3}{t^6(t+1)^3(t^2-t+1)^3(t-2)^6(t^2+2t+4)^6} \tag{2.14}$$

Therefore, solving for $\left((t^3+4)(t^3+6t^2+4)(t^6-6t^5+36t^4+8t^3-24t^2+16)\right)^3 = 1728 \cdot 4 \cdot f(t)^3$ and $t^6(t+1)^3(t^2-t+1)^3(t-2)^6(t^2+2t+4)^6 = 4f(t)^3 - 27g(t)^2$ gives:

$$\begin{aligned} f(t) &= \frac{1}{12 \cdot 2^{2/3}}\left(t^{12}+232t^9+960t^6+256t^3+256\right) \\ g(t) &= \frac{1}{216}\left(t^{18}-516t^{15}-12072t^{12}-24640t^9-30720t^6+6144t^3+4096\right) \end{aligned} \tag{2.15}$$

which we see to be polynomials of the correct degree. Note here that while $f = 0$ is solvable and has relatively simple roots, the corresponding equation for g is *not solvable.*

Now, the most general form of degree 12 and 18 polynomials in two variables are:

$$f(z_1, z_2) = \sum_{i=0}^{12} c_i z_1^i z_2^{12-i} \ , \quad g(z_1, z_2) = \sum_{i=0}^{18} c_i z_1^i z_2^{18-i} \ , \tag{2.16}$$

with complex coefficients c_i and d_i. We can readily find the coefficients c_i and d_i as well as constants a, b which allow (2.16) to be transformed into the form of (2.15). Setting, for example, We find that $a = 1$, $b = -2$, and that:

$$\begin{aligned} f(z_1,z_2) &= \tfrac{1}{8748 \cdot 2^{2/3}}\left(256z_1^{12} - 256z_2^3z_1^9 + 960z_2^6z_1^6 - 232z_2^9z_1^3 + z_2^{12}\right) \ ; \\ g(z_1,z_2) &= \tfrac{1}{4251528}\left(-4096z_1^{18} + 6144z_2^3z_1^{15} + 30720z_2^6z_1^{12} - 24640z_2^9z_1^9 + 12072z_2^{12}z_1^6 - 516z_2^{15}z_1^3 - z_2^{18}\right) \end{aligned} \tag{2.17}$$

We can readily find, using the same technique as above, the explicit Weierstraß equations, albeit with very complicated *algebraic* coefficients, of the remaining index 36 groups, as we had done for the index 24 cases in [3]. Though the complexity of the numerical expressions makes the presentation unworthwhile, it should be emphasized that all the coefficient we find for the Weierstraß models must be algebraic numbers. This is in accord with the deep theorem of Belyi that a rational map from a Riemann surface to $\mathbb{P}^1$ is Belyi (i.e., ramified only on 0, 1, ∞) if and only if there exists an algebraic equation for the Riemann surface over $\overline{\mathbb{Q}}$.

3 Semi-Stable Elliptic K3 Surfaces

We have mentioned above that the 9 index 24 subgroups are special in the sense that they correspond to semi-stable elliptic K3 surfaces. Indeed, exploring the physics of these was the subject of [3]. In this section, we will see how they belong to another important classification scheme which intersects nicely with our congruence subgroups.

In [10], the authors study semi-stable elliptic fibrations over $\mathbb{P}^1$ of K3 surfaces with six singular fibres. From the cusp widths, or, equivalently, the ramification data at ∞, in Table 1, we see that our subgroups all have 6 cusps and hence the index 24 cases belong to this class. In their paper, the authors give a complete list of possible fibre types for such fibrations - there are 112 such cases. Our nine index 24 subgroups of Γ discussed in the previous section correspond to nine of these 112 surfaces.

Remarkably, the elliptic j-invariants for all the 112 surfaces also have the property that, as rational maps from $\mathbb{P}^1$ to $\mathbb{P}^1$, they are Belyi. Furthermore, all the ramification indices at 0 are 3, and all the ramification indices at 1 are 2; that is, we have trivalent dessins! In [11], the dessins d'enfants for all 112 cases are constructed, along with explicit j-functions with coefficients in $\mathbb{Q}$ (For reference, we include the full list, emphasizing their clean, trivalent nature, in Appendix A). Those with coefficients in $\overline{\mathbb{Q}}\backslash\mathbb{Q}$ have yet to be computed, due to the complexity of finding exact roots to polynomials in high degree. This is thus an interesting open challenge. Now, because any finite trivalent graph is the Schreier coset graph of a finite index subgroup of the modular group, one naturally wonders whether we could attach a group to each of these dessins. Of course, only 9 of these will be congruence (they are all genus zero because the K3 surfaces are all fibred over $\mathbb{P}^1$), so what general subgroups can one encounter?

Indeed, we will see that just as those nine surfaces correspond to these congruence subgroups of Γ, it turns out that *all* the 112 surfaces will correspond to some (not necessarily congruence) subgroup of Γ. In this section we shall elaborate on the connections between these surfaces and the subgroups of Γ.

The structure of this section is as follows: first, we will discuss how the permutation representation of Γ on the right cosets of each modular subgroup defines a transitive permutation group known as a **cartographic group**. We then explain how these cartographic groups can be used to identify the generators of each of the modular subgroups (up to conjugacy), while also making clear the practical difficulties facing this approach. We then note that all the subgroups in question will be free groups on five generators, before proceeding to show how those generators can be identified computationally.

3.1 Permutation Representations and Cartographic Groups

Returning to our original discussion of the modular group in §2.1, let σ_0 and σ_1 denote the permutations induced by the respective actions of S and ST on the cosets of each subgroup. We can find a third permutation σ_∞ by imposing the following condition, thereby constructing a *permutation triple*:

$$\sigma_0 \cdot \sigma_1 \cdot \sigma_\infty = 1 \ . \tag{3.1}$$

Indeed, from the point of view of the dessins, there is an equivalent representation in terms of permutation triples [6, 7, 8]. Let d be the number of edges, labeled from 1 to d, and write elements of the symmetric group on d elements in the standard cycle notation as products over $(n_1, n_2, \ldots, n_k)$ where $n_i = 1, 2, \ldots, d$. Now, at the i-th black node write the labels for the incident edges clockwise, forming the cycle B_i; likewise, at the j-th white node write the labels for the incident edges counter-clockwise, forming the cycle W_j. Then, σ_0 is simply $\prod_i B_i$ while $\sigma_1 = \prod_j W_j$. Remarkably, σ_∞ as determined by (3.1) will have as many cycle products as there are faces, such that twice the length of each cycle is the number of edges of the corresponding face. Conversely, a *unique* dessin and Belyi map of degree d is thus determined.

As elements of the symmetric group, σ_0 and σ_1 can be easily computed from the Schreier coset graphs in [1] by following the procedure elaborated in [2], i.e., by noting that the doubly directed edges represent an element x of order 2, while the positively oriented triangles represent an element y of order 3. Since the graphs are connected, the group generated by x and y is transitive on the vertices. Clearly, σ_0 and σ_1 tell us which vertex of the coset graph is sent to which, i.e. which coset of the modular subgroup in question is sent to which by the action of Γ on the right cosets of this subgroup.

To express the cycle decomposition of the permutation σ_∞, we choose a vertex and apply x, then y, i.e., we take an edge (doubly oriented) followed by a side of a triangle (always in the positive orientation), until we return back to the original vertex. The length of this circuit gives the length of a cycle in σ_∞. In this way, we can construct the permutation decompositions σ_0, σ_1 and σ_∞ for every dessin under consideration. In turn, any two of these three permutation decompositions define a transitive permutation group. Of course, we only need two of these three permutations since the third will simply be the inverse of the product of the other two by (3.1), and we will henceforth focus on σ_0 and σ_1. These permutation groups are known as **cartographic groups**, and are discussed in e.g. [6]. We will shortly show how to extract the modular subgroup associated to a dessin from the cartographic group.

3.1.1 Cartographic Groups for the 112 K3 Surfaces

A priori, one would imagine that one could obtain the semi-stable elliptic K3 surfaces with 6 singular (I-type) singularities by partitioning 24 into 6 parts. There are 199 possibilities and [10] showed that only 112 are allowed. In general, one would try to partition 24 into at least 6 parts - the fact that there are at least 6 singular fibres is a restriction coming from the geometry of K3 surfaces. In our case, there are exactly 6 singular fibres, making these surfaces *extremal*. These surfaces are listed in [10].

We will thus distinguish our 112 extremal K3 surfaces by a 6-tuple partition $(I_1, \ldots, I_6)$ of 24; these will correspond to the I-type singular fibres as well as the ramification data (i.e., cusp widths) at infinity. A table of the cartographic groups for the partitions in [10] is given below. Now, dessins have intriguing number theoretic properties and the field of definition of (the coefficients in) the corresponding j-function as a Belyi map is an important quantity. For reference, we include the number field in brackets next to the partition. For simplicity, where we write $(\sqrt{-3})$, for example, we mean the field is $\mathbb{Q}(\sqrt{-3})$. Where it says e.g. "cubic", we mean it is $\mathbb{Q}$ extended by the root of some cubic polynomial.

We note that the partitions do not necessarily correspond uniquely to a K3 surface - when they do, the j-function is always defined over $\mathbb{Q}$. Where a single partition has multiple dessins defined over different number fields, we have for completeness computed the cartographic groups for all such cases, indicating explicitly the field as in [11]. We also keep the order of the entries as the same as the order of the dessins in [11] and in Appendix A.

To identify these cartographic groups, we use the corresponding dessins given in [11] to construct the Schreier coset graph in each case, following the procedure detailed in §2.3. We then read the permutations σ_0 and σ_1 directly off these graphs. As mentioned, together these permutations define a transitive permutation group, which is the cartographic group in question. Extensive use will be made of the computer software GAP [12]. The decomposition of each group in terms of better known groups can be identified using commands such as `StructureDescription()` and `TransitiveIdentification()` and it is these decompositions which are given in the ensuing table [†]. We follow [9] and use the following group notation: A (alternating), AL (affine linear), C (cyclic), E (elementary), S (symmetric). Direct products are denoted by "×", split extensions by ":", and wreath products by "≀". In addition, we use $L(n)$ to denote groups derived from linear groups as defined in Table 2 [9].

Where no easy decomposition is available, we have simply given the cartographic group's index in the GAP library of transitive permutation groups (here a number of the form t24n...).

[†] We thank Alexander Hulpke for helping to identify many of the decompositions.

Name	**Size**	**Type**
$L(6)$	60	$PSL(2,5) = A_5(6)$
$L(7)$	168	$PSL(3,2)$
$L(8)$	168	$PSL(2,7)$
$L(9)$	504	$PSL(2,8)$
$L(10)$	360	$PSL(2,9)$
$L(11)$	660	$PSL(2,11)(11)$
$L(12)$	660	$PSL(2,11)$
$L(13)$	5616	$PSL(3,3)$
$L(14)$	1092	$PSL(2,13)$
$L(15)$	20160	$PSL(4,2) = A_8(15)$

Table 2: *The permutation groups* $L(n)$

The number after ‘t’ in this index gives the degree transitive permutation group; the number after the ‘n’ gives the location that group in the GAP library of transitive permutation groups for that particular group degree.

Partition (K3 Surface)	**Cartographic Group**	**Cartographic Group Order**
$19,1,1,1,1,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$18,2,1,1,1,1$ $(\mathbb{Q})$	t24n18368 = t24n16559 $\times C_2$	$2^{11} \cdot 3^4$
$18,2,1,1,1,1$ $(\sqrt{-3})$	A_{24}	$24!/2$
$18,2,1,1,1,1$ $(\sqrt{-3})$	A_{24}	$24!/2$
$17,3,1,1,1,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$17,2,2,1,1,1$ $\left(\sqrt{17}\right)$	A_{24}	$24!/2$
$17,2,2,1,1,1$ $\left(\sqrt{17}\right)$	A_{24}	$24!/2$
$16,4,1,1,1,1$ $(\mathbb{Q})$	$(((C_8 \times C_2) : C_8) : C_3) : C_2$	$2^8 \cdot 3$
$16,3,2,1,1,1$ $(\mathbb{Q})$	t24n24710 = 2^{12} : t18n762	$2^{17} \cdot 3^7$
$16,3,2,1,1,1$ $(\sqrt{-2})$	A_{24}	$24!/2$
$16,3,2,1,1,1$ $(\sqrt{-2})$	A_{24}	$24!/2$
$16,2,2,2,1,1$ $(\mathbb{Q})$	$((C_8 \times C_8) : C_3) : C_2$	$2^7 \cdot 3$
$15,5,1,1,1,1$ $(\sqrt{-15})$	A_{24}	$24!/2$
$15,5,1,1,1,1$ $(\sqrt{-15})$	A_{24}	$24!/2$
$15,4,2,1,1,1$ $(\sqrt{-15})$	A_{24}	$24!/2$
$15,4,2,1,1,1$ $(\sqrt{-15})$	A_{24}	$24!/2$
$15,3,3,1,1,1$ $(\mathbb{Q})$	$A_6 \wr A_4$	$2^{14} \cdot 3^9 \cdot 5^4$

$15, 3, 2, 2, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$15, 3, 2, 2, 1, 1$ $\left(\sqrt{-15}\right)$	A_{24}	$24!/2$
$15, 3, 2, 2, 1, 1$ $\left(\sqrt{-15}\right)$	A_{24}	$24!/2$
$14, 6, 1, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$14, 6, 1, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$14, 5, 2, 1, 1, 1$ $(\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$14, 5, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$14, 5, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$14, 5, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$14, 4, 3, 1, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$14, 4, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$	$(E_8 : L(7)) \wr S_3 = \mathrm{ASL}(3, 2) \wr S_3$	$2^{19} \cdot 3^4 \cdot 7^3$
$14, 4, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$	$(E_8 : L(7)) \wr S_3 = \mathrm{ASL}(3, 2) \wr S_3$	$2^{19} \cdot 3^4 \cdot 7^3$
$14, 3, 3, 2, 1, 1$ $(\mathbb{Q})$	$L(8) \wr S_3 = \mathrm{PSL}(2, 7) \wr S_3$	$2^{10} \cdot 3^4 \cdot 7^3$
$14, 3, 3, 2, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$14, 3, 2, 2, 2, 1$ $(\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$13, 7, 1, 1, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$13, 6, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$13, 6, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$13, 5, 3, 1, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$13, 5, 2, 2, 1, 1$ $\left(\sqrt{65}\right)$	A_{24}	$24!/2$
$13, 5, 2, 2, 1, 1$ $\left(\sqrt{65}\right)$	A_{24}	$24!/2$
$13, 4, 3, 2, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$13, 3, 3, 2, 2, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$12, 7, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$12, 7, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$	A_{24}	$24!/2$
$12, 6, 3, 1, 1, 1$ $(\mathbb{Q})$	$A_6 \wr A_4$	$2^{14} \cdot 3^9 \cdot 5^4$
$12, 6, 2, 2, 1, 1$ $(\mathbb{Q})$	t24n12260	$2^{11} \cdot 3^2$
$12, 6, 2, 2, 1, 1$ $(\mathbb{Q})$	$(L(8) \wr S_3) \cap A_{24}$	$2^{12} \cdot 3^4 \cdot 7^3$
$12, 5, 3, 2, 1, 1$ (quartic)	A_{24}	$24!/2$
$12, 5, 3, 2, 1, 1$ (quartic)	A_{24}	$24!/2$

$12, 5, 3, 2, 1, 1$ (quartic)	A_{24}	$24!/2$
$12, 5, 3, 2, 1, 1$ (quartic)	A_{24}	$24!/2$
$12, 5, 2, 2, 2, 1$ $(\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$12, 4, 4, 2, 1, 1$ $(\mathbb{Q})$	t24n21388 $= 2^{12} \wr$ t18n97	$2^{15} \cdot 3^3$
$12, 4, 3, 3, 1, 1$ $(\mathbb{Q})$	$S_4 \times A_4$	$4! \cdot 4!/2$
$12, 4, 3, 3, 1, 1$ $(\mathbb{Q})$	$(\mathrm{PSL}\,(2, 5) \wr A_4) : C_2$	$2^{11} \cdot 3^5 \cdot 5^4$
$12, 4, 3, 2, 2, 1$ $(\mathbb{Q})$	t24n23732 $= 2^{12} \wr$ t18n346	$2^{15} \cdot 3^5$
$12, 4, 3, 2, 2, 1$ $\left(\sqrt{-3}\right)$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$12, 4, 3, 2, 2, 1$ $\left(\sqrt{-3}\right)$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$12, 3, 3, 3, 2, 1$ $(\mathbb{Q})$	$(A_6 \wr A_4) : C_2$	$2^{15} \cdot 3^9 \cdot 5^4$
$12, 3, 3, 2, 2, 2$ $(\mathbb{Q})$	t24n12261	$2^{11} \cdot 3^2$
$11, 9, 1, 1, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$11, 8, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 8, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 8, 2, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 7, 3, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 7, 3, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 7, 3, 1, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 7, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$	A_{24}	$24!/2$
$11, 7, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$	A_{24}	$24!/2$
$11, 6, 4, 1, 1, 1$ $\left(\sqrt{33}\right)$	A_{24}	$24!/2$
$11, 6, 4, 1, 1, 1$ $\left(\sqrt{33}\right)$	A_{24}	$24!/2$
$11, 6, 3, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 6, 3, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 6, 3, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 5, 5, 1, 1, 1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$11, 5, 4, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 5, 4, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 5, 4, 2, 1, 1$ (cubic)	A_{24}	$24!/2$
$11, 5, 3, 3, 1, 1$ $\left(\sqrt{5}\right)$	A_{24}	$24!/2$

$11,5,3,3,1,1\ \left(\sqrt{5}\right)$	A_{24}	$24!/2$
$11,5,3,2,2,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$11,4,4,3,1,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$11,4,3,3,2,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$10,10,1,1,1,1\ (\mathbb{Q})$	$((C_2 \times C_2 \times C_2 \times C_2 \times C_2) : A_5) : C_2$	$2^8 \cdot 3 \cdot 5$
$10,10,1,1,1,1\ \left(\sqrt{5}\right)$	t24n22755	$2^{14} \cdot 3 \cdot 5 \cdot 11$
$10,10,1,1,1,1\ \left(\sqrt{5}\right)$	t24n22755	$2^{14} \cdot 3 \cdot 5 \cdot 11$
$10,9,2,1,1,1\ \left(\sqrt{5}\right)$	A_{24}	$24!/2$
$10,9,2,1,1,1\ \left(\sqrt{5}\right)$	A_{24}	$24!/2$
$10,8,3,1,1,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$10,7,4,1,1,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$10,7,3,2,1,1\ \left(\sqrt{21}\right)$	A_{24}	$24!/2$
$10,7,3,2,1,1\ \left(\sqrt{21}\right)$	A_{24}	$24!/2$
$10,7,2,2,2,1\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,6,5,1,1,1\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,6,5,1,1,1$ (cubic)	A_{24}	$24!/2$
$10,6,5,1,1,1$ (cubic)	A_{24}	$24!/2$
$10,6,5,1,1,1$ (cubic)	A_{24}	$24!/2$
$10,6,4,2,1,1\ (\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,6,4,2,1,1\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,6,3,2,2,1\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,5,5,2,1,1\ \left(\sqrt{5}\right)$	$((C_2 \times C_2 \times C_2 \times C_2 \times C_2) : A_5) : C_2$	$2^8 \cdot 3 \cdot 5$
$10,5,5,2,1,1\ \left(\sqrt{5}\right)$	$((C_2 \times C_2 \times C_2 \times C_2 \times C_2) : A_5) : C_2$	$2^8 \cdot 3 \cdot 5$
$10,5,4,2,2,1$ (cubic)	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,5,4,2,2,1$ (cubic)	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,5,4,2,2,1$ (cubic)	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,5,3,2,2,2\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,4,4,3,2,1\ (\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$10,4,3,3,2,2\ (\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$9,9,2,2,1,1\ (\mathbb{Q})$	t24n16559	$2^{10} \cdot 3^4$

$9,9,2,2,1,1$ (cubic)	t24n24970	$2^{21}\cdot 3^5\cdot 5^2\cdot 7\cdot 11$
$9,9,2,2,1,1$ (cubic)	t24n24970	$2^{21}\cdot 3^5\cdot 5^2\cdot 7\cdot 11$
$9,9,2,2,1,1$ (cubic)	t24n24970	$2^{21}\cdot 3^5\cdot 5^2\cdot 7\cdot 11$
$9,8,3,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,8,3,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,8,3,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,7,5,1,1,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$9,7,4,2,1,1$ $(\sqrt{-7})$	A_{24}	$24!/2$
$9,7,4,2,1,1$ $(\sqrt{-7})$	A_{24}	$24!/2$
$9,7,3,2,2,1$ (cubic)	A_{24}	$24!/2$
$9,7,3,2,2,1$ (cubic)	A_{24}	$24!/2$
$9,7,3,2,2,1$ (cubic)	A_{24}	$24!/2$
$9,6,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,6,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,6,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$9,6,4,3,1,1$ $(\mathbb{Q})$	$(A_6 \wr A_4) : C_2$	$2^{15}\cdot 3^9\cdot 5^4$
$9,6,3,3,2,1$ (cubic)	$(A_6 \wr A_4) : C_2$	$2^{15}\cdot 3^9\cdot 5^4$
$9,6,3,3,2,1$ (cubic)	$(A_6 \wr A_4) : C_2$	$2^{15}\cdot 3^9\cdot 5^4$
$9,6,3,3,2,1$ (cubic)	$(A_6 \wr A_4) : C_2$	$2^{15}\cdot 3^9\cdot 5^4$
$9,5,5,2,2,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$9,5,4,3,2,1$ (cubic)	A_{24}	$24!/2$
$9,5,4,3,2,1$ (cubic)	A_{24}	$24!/2$
$9,5,4,3,2,1$ (cubic)	A_{24}	$24!/2$
$9,5,3,3,3,1$ $(\mathbb{Q})$	$A_6 \wr A_4$	$2^{14}\cdot 3^9\cdot 5^4$
$9,4,3,3,3,2$ $(\mathbb{Q})$	$A_6 \wr A_4$	$2^{14}\cdot 3^9\cdot 5^4$
$8,8,4,2,1,1$ $(\mathbb{Q})$	$(((C_4\times C_2):C_4):C_3):C_2$	$2^6\cdot 3$
$8,8,3,3,1,1$ $(\mathbb{Q})$	t24n14598 (Index 2 in t8n14 $\wr S_3$)	$2^9\cdot 3^4$
$8,8,3,3,1,1$ $(\sqrt{-2})$	t24n24970	$2^{21}\cdot 3^5\cdot 5^2\cdot 7\cdot 11$
$8,8,3,3,1,1$ $(\sqrt{-2})$	t24n24970	$2^{21}\cdot 3^5\cdot 5^2\cdot 7\cdot 11$
$8,8,2,2,2,2$ $(\mathbb{Q})$	$((C_4\times C_4):C_3):C_2$	$2^5\cdot 3$

$8,7,6,1,1,1\ (\sqrt{-3})$	A_{24}	$24!/2$
$8,7,6,1,1,1\ (\sqrt{-3})$	A_{24}	$24!/2$
$8,7,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$8,7,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$8,7,5,2,1,1$ (cubic)	A_{24}	$24!/2$
$8,7,4,3,1,1\ (\sqrt{-6})$	A_{24}	$24!/2$
$8,7,4,3,1,1\ (\sqrt{-6})$	A_{24}	$24!/2$
$8,7,4,2,2,1\ (\sqrt{-7})$	$(E_8 : L(7)) \wr S_3 = \mathrm{ASL}(3,2) \wr S_3$	$2^{19} \cdot 3^4 \cdot 7^3$
$8,7,4,2,2,1\ (\sqrt{-7})$	$(E_8 : L(7)) \wr S_3 = \mathrm{ASL}(3,2) \wr S_3$	$2^{19} \cdot 3^4 \cdot 7^3$
$8,7,3,3,2,1\ (\mathbb{Q})$	A_{24}	$24!/2$
$8,6,6,2,1,1\ (\sqrt{-3})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,6,6,2,1,1\ (\sqrt{-3})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,6,5,3,1,1\ \left(\sqrt{5}\right)$	A_{24}	$24!/2$
$8,6,5,3,1,1\ \left(\sqrt{5}\right)$	A_{24}	$24!/2$
$8,6,5,2,2,1\ (\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,6,4,3,2,1\ \left(\sqrt{2}\right)$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,6,4,3,2,1\ \left(\sqrt{2}\right)$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,6,4,2,2,2\ (\mathbb{Q})$	$\mathrm{t24n22747} = 2^{12} \wr \mathrm{t18n217}$	$2^{15} \cdot 3^4$
$8,5,4,3,3,1\ \left(\sqrt{10}\right)$	A_{24}	$24!/2$
$8,5,4,3,3,1\ \left(\sqrt{10}\right)$	A_{24}	$24!/2$
$8,5,4,3,2,2\ (\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$8,4,4,4,3,1\ (\mathbb{Q})$	$\mathrm{t24n24710} = 2^{12} : \mathrm{t18n762}$	$2^{17} \cdot 3^7$
$8,4,4,4,2,2\ (\mathbb{Q})$	$(((C_4 \times C_2) : C_4) : C_3) : C_2$	$2^6 \cdot 3$
$7,7,7,1,1,1\ (\mathbb{Q})$	PSL(3,2)	$2^3 \cdot 3 \cdot 7$
$7,7,5,3,1,1\ \left(\sqrt{21}\right)$	A_{24}	$24!/2$
$7,7,5,3,1,1\ \left(\sqrt{21}\right)$	A_{24}	$24!/2$
$7,7,4,4,1,1\ (\sqrt{-7})$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$7,7,4,4,1,1\ (\sqrt{-7})$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$7,7,3,3,2,2\ \left(\sqrt{7}\right)$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$7,7,3,3,2,2\ \left(\sqrt{7}\right)$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$

$7,6,6,2,2,1$ $(\mathbb{Q})$	$L(8) \wr S_3 = \mathrm{PSL}(2,7) \wr S_3$	$2^{10} \cdot 3^4 \cdot 7^3$
$7,6,5,4,1,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$7,6,5,3,2,1$ (cubic)	A_{24}	$24!/2$
$7,6,5,3,2,1$ (cubic)	A_{24}	$24!/2$
$7,6,5,3,2,1$ (cubic)	A_{24}	$24!/2$
$7,6,4,4,2,1$ $(\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$7,5,5,4,2,1$ $\left(\sqrt{2}\right)$	A_{24}	$24!/2$
$7,5,5,4,2,1$ $\left(\sqrt{2}\right)$	A_{24}	$24!/2$
$7,5,5,3,2,2$ $(\mathbb{Q})$	A_{24}	$24!/2$
$7,5,4,4,3,1$ $(\mathbb{Q})$	A_{24}	$24!/2$
$7,5,4,3,3,2$ $(\mathbb{Q})$	A_{24}	$24!/2$
$6,6,6,4,1,1$ $(\mathbb{Q})$	t24n12261	$2^{11} \cdot 3^2$
$6,6,6,2,2,2$ $(\mathbb{Q})$	$A_4 \times S_3$	72
$6,6,5,5,1,1$ $\left(\sqrt{3}\right)$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$6,6,5,5,1,1$ $\left(\sqrt{3}\right)$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$6,6,5,4,2,1$ $(\mathbb{Q})$	$(S_8 \wr S_3) \cap A_{24}$	$2^{21} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$6,6,5,3,3,1$ $(\mathbb{Q})$	$L(6) \wr A_4 = \mathrm{PSL}(2,5) \wr A_4$	$2^{10} \cdot 3^5 \cdot 5^4$
$6,6,4,4,2,2$ $(\mathbb{Q})$	$(((C_2 \times C_2\,((C_2 \times C_2 \times C_2 \times C_2) : C_3)) : C_3) : C_3) : C_2$	$2^7 \cdot 3^3$
$6,6,4,3,3,2$ $(\mathbb{Q})$	t24n12260	$2^{11} \cdot 3^2$
$6,6,3,3,3,3$ $(\mathbb{Q})$	$((C_4 \times C_4) : C_3) : C_2$	$2^5 \cdot 3$
$6,5,4,4,3,2$ $(\mathbb{Q})$	$A_8 \wr S_3$	$2^{19} \cdot 3^7 \cdot 5^3 \cdot 7^3$
$5,5,5,5,2,2$ $(\mathbb{Q})$	$((C_2 \times C_2 \times C_2 \times C_2 \times C_2) : A_5) : C_2$	$2^8 \cdot 3 \cdot 5$
$5,5,4,4,3,3$ $(\mathbb{Q})$	t24n24970	$2^{21} \cdot 3^5 \cdot 5^2 \cdot 7 \cdot 11$
$4,4,4,4,4,4$ $(\mathbb{Q})$	S_4	$4!$

3.2 Modular Subgroup Generators from Cartographic Groups

Having obtained the cartographic groups from the dessins, we can now proceed to extract the corresponding modular subgroup. The permutations σ_0, σ_1 and σ_∞ give the permutation representations of Γ on the right cosets of each modular subgroup in question. This can be confirmed by computing the σ_i where the subgroups are known; for example, we know the nine index 24 congruence subgroups discussed in §2.2 and in [3], so we can obtain the σ_i

directly to check. We again make use [12, 13] of GAP or Sage and find that such permutation data matches that obtained from the Schreier coset graphs.

At this point, let us extend our terminology and discuss how to extract in general the generators of a subgroup H. We are given G (in our case, G is the modular group) and the permutation representation of G acting on the right cosets of H defining a cartographic group C. Knowing G and C but not H, it is possible to identify the generators of H, up to conjugacy. The procedure is as follows:

PROPOSITION 3 *Choose a set of generators for G. Draw the Cayley graph of the action of G on the cosets $G\backslash H$ (i.e., the Cayley graph of the cartographic group C). Choose a set of generators for the fundamental group of the graph. For each generator of the fundamental group, multiply the generators of G corresponding to the edges together in order. This will be a set of generators for H.*

Proof: [‡] Each element of H is a product of the generators of G which, in the action of G on $G\backslash H$, preserves the identity coset. Thus each element of H is a product of generators that forms a cycle of the Cayley graph. Following the homotopy of that cycle to a composition of the generators of the fundamental group does not change the element it refers to. Therefore, we now have an expression for each element of H in terms of the supposed generators of H, so they are in fact the generators of H.

In principle then, we have found that to identify the generators of H, we need only consider the Cayley graph for the corresponding C. The cycles of the Cayley graph give the generators of H. There are, however, two significant drawbacks to this method:

1. This solution assumes that we know a fixed element of $G\backslash H$ which can be used as the basepoint for the fundamental group, namely the identity coset H. If we do not know this (as is the case we are presently confronting), we cannot recover H exactly, but only up to conjugacy, since the permutation actions of G on $G\backslash H$ and $G\backslash H'$ are permutationally isomorphic if and only if H and H' are conjugate subgroups. As noted, many of the C in our case are A_{24}. This means that this method will only take us so far in identifying the generators of the H.

2. There is a practical problem in implementing this method. The reason for this is due to the fact that if a directed graph has v vertices and e edges, then there are $2^{(e-v+1)} - 1$ cycles. Now, the C of lowest order in the table in §3.1.1 is S_4, with order 24, so its Cayley graph will have $v = 24$ and $e \geq 24$. In fact though, the orders of the majority of the C are substantially higher; the order of A_{24}, for example, is

---

[‡] We thank Will Sawin for this argument.

$24!/2 = 310224200866619719680000$. Hence, to enumerate all these cycles would require unrealistic computational power; and even then, there will be a huge redundancy in the generators of H obtained from these cycles.

As a result of these points, it turns out that this method of identifying the generators of the H from the G and C in each case is not ideal. For this reason, we now consider alternative procedures.

3.3 Subgroup Presentations: Free Groups on Five Generators

It turns out that there is a so-called **Reidemeister-Schreier process** (discussed at length in [14]) which enables one to compute presentations of subgroups H of finite index in a group G defined by finite presentations, taking as input either the σ_0, σ_1 and σ_∞, or the Schreier coset graphs, or standardized coset tables (which we recall is just a table listing how each element of G sends one coset to another). All these three alternative inputs are, of course, equivalent.

We can use this process to find the presentations of all the modular subgroups of interest. In our case, G is $\Gamma = PSL(2,\mathbb{Z})$, the presentation of which is $\langle X|R\rangle = \langle x,y|x^2,y^3\rangle$, where X is the set of the generators of G, and R is the set of relations between these generators.

Step I: In brief, the Reidemeister-Schreier process defines a set Y of Schreier generators[§] y_{tx} of H and computes a word $P[\alpha,x]$ in $Y \cup Y^{-1}$ (i.e. either one of the generators in Y or its inverse) for each $\alpha \in \Omega$ where Ω is the set of cosets of H and $x \in (X \cup X^{-1})$ where X is the set of generators of G. Each $P[\alpha,x]$ is either the empty word, or a Schreier generator in Y (when $x \in X$), or an inverse of a Schreier generator (when $x \in X^{-1}$). We regard Y and P as being new components of the coset table $\mathcal{C}$ of H, and define a *modified coset table* $\mathcal{C}_M$ to be an ordinary coset table endowed with these two extra components. In practice, what this amounts to is a word $P[\alpha,x]$ being placed in every cell of $\mathcal{C}$; the purpose of this shall be explained presently. The first part of the Reidemeister-Schreier algorithm (given in e.g. [14]) produces this $\mathcal{C}_M$.

[§] In this footnote we recall the definition of a Schreier generator. Let G be a group generated by a set X, and let $A = X \cup X^{-1}$. Then $G \cong F/N$ for some N, where $F = F_X$ is the free group on X, and we shall assume that $G = F/N$. Let $H = E/N$ be a subgroup of G, and let T be a set of reduced words in A^* that form a right transversal of E in F. We shall always assume that T contains the empty word ε as the representative of the coset E. Hence the images of T in G form a right transversal of H in G. For a word $w \in A^*$, we denote the unique element of $T \cap Ew$ by $\overline{w}$.

Now, let $w = x_1x_2\cdots x_r \in A^*$ and, for $0 \le i \le r$, let $t_i := \overline{x_1x_2\cdots x_i}$, where $t_0 = \varepsilon$. Then we have $w =_F \left(t_0x_1t_1^{-1}\right)\left(t_1x_2t_2^{-1}\right)\cdots\left(t_{r-1}x_rt_r^{-1}\right)t_r$. Define the subset $\hat{Y}$ of E by $\hat{Y} := \left\{tx\overline{tx}^{-1} | t \in T, x \in X, tx \neq_F \overline{tx}^{-1}\right\}$. Then, the theorem of Schreier tells us that the subgroup E of F is generated by $\hat{Y}$. Similarly the subgroup H of G is generated by the images of $\hat{Y}$ in G. The set $\hat{Y}$ is called the set of *Schreier generators* of E with respect to the group F. We let Y be a set of new symbols corresponding to $\hat{Y}$, where we denote the element of Y corresponding to $tx\overline{tx}^{-1} \in Y$ by y_{tx}. For further discussion, see [14].

Let us give an example for the partition $4, 4, 4, 4, 4, 4$ - from [3], we know this corresponds to the principal congruence subgroup $\Gamma(4)$. Using the permutation data given by the Schreier coset graph (or, alternatively, given directly in [10]), we can construct a standardized coset table $\mathcal{C}$ for this modular subgroup (for details of the standardization procedure for coset tables, see [14]). The Reidemeister-Schreier process then gives us the modified coset table $\mathcal{C}_M$ for this case.

For $\Gamma(4)$, this modified coset table is given in Table 4, where columns label the generators of G and their inverses (i.e. the elements of $X \cup X^{-1}$), and rows label the cosets of H (i.e. the elements of Ω). The ramification structure of all the entries in [11] tells us that all the modular subgroups of interest will be index 24 in Γ (using the results of §2.2), and so will have 24 right cosets.

Step II: After this modified coset table $\mathcal{C}_M$ is produced, the second half of the Reidemeister-Schreier algorithm outputs the presentation of the H in question. Having already produced the set of Schreier generators Y for H, the only thing left to do is to compute the relations between these generators. This is done as follows: let α^x be the number in the αth row, xth column of the coset table. Let $\beta := \alpha^x$, β is the coset to which α is sent by the action of the generator x. If that entry in the table also contains a word $P[\alpha, x]$, record that word. Then repeat the process for β, x, updating β so that $\beta := \beta^x$, and appending any additional words $P[\beta, x]$ to the previously recorded words (if any). When we return to $\beta = \alpha$, we have completed a full "cycle", and can stop. The word we have written down will be one of the relators for H. Repeat this process for all $\alpha \in \Omega$ and $x \in X$ to find all the relators of H.

Geometric Interpretation: Interestingly, the whole Reidemeister-Schreier process can be formulated geometrically. This formulation is as follows (see [15] for more detail): to find a presentation for a subgroup H of G, choose a spanning-tree in the associated coset graph, label the unused edges with (irredundant) Schreier generators, and then apply the relators in R to each of the vertices in turn to obtain the relators of H. In fact, deploying this form of the Reidemeister-Schreier process saves significant time in our case, since we can work directly with the coset graphs obtained from the dessins, rather than having to compute the standardized coset tables in every case.

In any case, running the second part of the Reidemeister-Schreier algorithm enables us to compute the presentation for the H in question. The result for $\Gamma(4)$ is given below, where

	x		x^{-1}		y		y^{-1}	
1		2	a^{-1}	2		3		4
2	a	1		1		5		6
3		7	d^{-1}	7	b	4		1
4		8	e^{-1}	8		1	b^{-1}	3
5		9	f^{-1}	9	c	6		2
6		10	g^{-1}	10		2	c^{-1}	5
7	d	3		3		11		12
8	e	4		4		13		14
9	f	5		5		15		16
10	g	6		6		17		18
11	h	18	o^{-1}	18	i	12		7
12		19	p^{-1}	19		7	i^{-1}	11
13		20	q^{-1}	20	j	14		8
14	k	15	l^{-1}	15		8	j^{-1}	13
15	l	14	k^{-1}	14	m	16		9
16		21	s^{-1}	21		9	m^{-1}	15
17		22	t^{-1}	22	n	18		10
18	o	11	h^{-1}	11		10	n^{-1}	17
19	p	12		12		23	r^{-1}	20
20	q	13		13	r	19	w^{-1}	23
21	s	16		16		24	u^{-1}	22
22	t	17		17	u	21	y^{-1}	24
23	v	24	x^{-1}	24	w	20		19
24	x	23	v^{-1}	23	y	22		21

Table 4: *The modified coset table $\mathcal{C}_M$ for $\Gamma(4)$. Columns indicate the generators of G and their inverses; rows indicate the 24 right cosets of H. Numbers in the table indicate the action of G on the right cosets of H. The words in the table are the associated Schreier generators of H and their inverses. The ordinary coset table $\mathcal{C}$, by comparison, contains no such words.*

we have omitted redundant cyclic conjugates of relators:

$$\langle [a..y] \,| a, d, e, f, g, ho, p, q, kl, s, t, vx, b, c, i, j, m, n, wr, yu \rangle \tag{3.2}$$

This result can be further simplified by a series of Tietze transformations [14] as follows.

First, all generators with trivial monadic relators (e.g. $a, d, e, \ldots$ etc.) can be omitted. Second, one of the two generators in each of the "double" relators (e.g. ho for h or o, kl for k or l, etc.) can be omitted, since the one will simply be the inverse of the other. Making these simplifications gives the following final presentation for $\Gamma(4)$:

$$\langle h, k, w, x, y | \rangle \tag{3.3}$$

This tells us that $\Gamma(4)$ is a free group on 5 generators. Ultimately this should not be surprising, since the modular curve $X(4)$ is $\mathbb{P}^1$ with six cusps and no elliptic points, so its fundamental group is the free group on 5 generators. Indeed, since this is true of all the modular curves corresponding to the entires in [11], it should be equally unsurprising to find that the Reidemeister-Schreier process returns (3.3) in every case. Still, this is certainly a step in the right direction on our journey to identifying all the modular subgroups of interest, since we have shown explicitly that we are searching for free groups on five generators in every case. If we can find these five generators for every group as explicit 2×2 matrices, our task will be complete.

3.4 Generators for the Modular Subgroups

We can, in fact, find the generators [¶] for all the subgroups of interest using GAP. It is testament to the striking wealth of structure encoded in Γ that all these free groups will turn out to be distinct, with no two being isomorphic to one another.

First, we use the permutation data σ_0, σ_1 obtained from each of the Schreier coset graphs (in turn obtained from each of the dessins) to find the group homomorphism by images between G and H. We then use this to define the H in question. Finally, we use the GAP command `IsomorphismFpGroup(G)`, which returns an isomorphism from the given H to a finitely presented group isomorphic to H. This function first chooses a set of generators of H and then computes a presentation in terms of these generators.

To give an example, consider again the partition 4, 4, 4, 4, 4, 4 - which corresponds to the congruence subgroup $\Gamma(4)$. We can find a set of generators as 2×2 matrices for this group by implementing the following code in GAP:

```
gap> f:=FreeGroup("x","y");
<free group on the generators [ x, y ]>
gap>gamma:=f/[f.1^2,f.2^3];
<fp group on the generators [ x, y ]>
gap> hom:=GroupHomomorphismByImages(gamma,Group(sigma_0,sigma_1),
```

[¶] We thanks Alexander Hulpke for helpful guidance here.

```
>GeneratorsOfGroup(gamma),[sigma_0,sigma_1]);
[x, y] ->
[(1,10)(2,4)(3,24)(5,7)(6,21)(8,12)(9,18)(11,14)(13,16)(15,23)
(17,19)(20,22), (1,2,3)(4,5,6)(7,8,9)(10,11,12)(13,14,15)
(16,17,18)(19,20,21)(22,23,24)]
gap> gamma4:=PreImage(hom,Stabilizer(Image(hom),1));
Group(<fp, no generators known>)
gap> iso:=IsomorphismFpGroup(gamma4);
[ <[ [ 1, 1 ] ]|y*x*y*x*y*x^-1*y*x^-1>,
<[ [ 2, 1 ] ]|y^-1*x*y^-1*x*y^-1*x^ -1*y^-1*x^-1>,
<[ [ 3, 1 ] ]|y*x*y^-1*x*y^-1*x^-1*y^-1*x^-1*y>,
<[ [ 4, 1 ] ]|x*y*x*y^-1*x*y^-1*x^-1*y^-1*x^-1*y*x^-1>,
<[ [ 5, 1 ] ]|y*x*y^-1*x*y*x*y^-1*x^-1*y*x^-1*y^-1*x^-1> ]
-> [ F1, F2, F3, F4, F5 ]
```

Here, the only input in each case are the permutation data σ_0, σ_1. Once the output has been obtained, we will see the five generators $[F1, F2, F3, F4, F5]$, as indicated in the last line of the output, as functions of x and y, i.e. the generators of Γ. Now, returning to the matrices (2.4), the only thing left to do is to multiply together the matrices and their inverses as indicated. This will produce the generators of each H, as desired.

3.4.1 Catalogue of Generators as Explicit 2×2 Matrices

In this section we give the generators of each of the modular subgroups corresponding to the entries in Table at the end of §3.1.1. For each of the 112 K3 surfaces, denoted as a 6-tuple partition of 24 (the notation is as before, we also include the number field of definition for the associated j-function as Belyi map), we write the five matrix generators explicitly.

Partition	**Generators**
$19,1,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix} \begin{pmatrix} 18 & -5 \\ 25 & 0 \end{pmatrix} \begin{pmatrix} 20 & 9 \\ -49 & -22 \end{pmatrix}$
$18,2,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} -5 & -2 \\ 18 & 7 \end{pmatrix} \begin{pmatrix} 5 & 1 \\ -36 & -7 \end{pmatrix} \begin{pmatrix} 17 & 4 \\ -81 & -19 \end{pmatrix}$
$18,2,1,1,1,1$ $\left(\sqrt{-3}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix} \begin{pmatrix} -19 & -8 \\ 50 & 21 \end{pmatrix} \begin{pmatrix} 43 & 16 \\ -121 & -45 \end{pmatrix}$
$18,2,1,1,1,1$ $\left(\sqrt{-3}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} 9 & 4 \\ -25 & -11 \end{pmatrix} \begin{pmatrix} -7 & -2 \\ 32 & 9 \end{pmatrix} \begin{pmatrix} 5 & 1 \\ -36 & -7 \end{pmatrix}$
$17,3,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} -8 & -3 \\ 27 & 10 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ -25 & -6 \end{pmatrix} \begin{pmatrix} 20 & 9 \\ -49 & -22 \end{pmatrix}$
$17,2,2,1,1,1$ $\left(\sqrt{17}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix} \begin{pmatrix} -7 & -2 \\ 32 & 9 \end{pmatrix} \begin{pmatrix} 6 & 1 \\ -49 & -8 \end{pmatrix} \begin{pmatrix} 21 & 4 \\ -121 & -23 \end{pmatrix}$

$17,2,2,1,1,1\ \left(\sqrt{17}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}-27&-8\\98&29\end{pmatrix}\begin{pmatrix}59&16\\-225&-61\end{pmatrix}$
$16,4,1,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}7&4\\-16&-9\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}11&9\\-16&-13\end{pmatrix}\begin{pmatrix}23&9\\-64&-25\end{pmatrix}$
$16,3,2,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-11&-3\\48&13\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}\begin{pmatrix}29&9\\-100&-31\end{pmatrix}$
$16,3,2,1,1,1\ \left(\sqrt{-2}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}9&4\\-25&-11\end{pmatrix}\begin{pmatrix}-3&-1\\22&7\end{pmatrix}\begin{pmatrix}-11&-3\\48&13\end{pmatrix}$
$16,3,2,1,1,1\ \left(\sqrt{-2}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}-13&-6\\37&17\end{pmatrix}\begin{pmatrix}-29&-12\\75&31\end{pmatrix}$
$16,2,2,2,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}-11&-2\\72&13\end{pmatrix}\begin{pmatrix}7&1\\-64&-9\end{pmatrix}$
$15,5,1,1,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}9&4\\-25&-11\end{pmatrix}\begin{pmatrix}3&1\\-19&-6\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}$
$15,5,1,1,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}11&5\\-31&-14\end{pmatrix}\begin{pmatrix}64&25\\-169&-66\end{pmatrix}$
$15,4,2,1,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}3&2\\-14&-9\end{pmatrix}\begin{pmatrix}-5&-2\\18&7\end{pmatrix}\begin{pmatrix}34&25\\-49&-36\end{pmatrix}\begin{pmatrix}19&16\\-25&-21\end{pmatrix}$
$15,4,2,1,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-14&-11\end{pmatrix}\begin{pmatrix}7&4\\-16&-9\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}13&4\\-49&-15\end{pmatrix}$
$15,3,3,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}4&3\\-15&-11\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}$
$15,3,2,2,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}2&1\\-11&-5\end{pmatrix}\begin{pmatrix}-19&-6\\73&23\end{pmatrix}\begin{pmatrix}-41&-12\\147&43\end{pmatrix}$
$15,3,2,2,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}13&4\\-49&-15\end{pmatrix}\begin{pmatrix}-4&-1\\33&8\end{pmatrix}\begin{pmatrix}-14&-3\\75&16\end{pmatrix}$
$15,3,2,2,1,1\ \left(\sqrt{-15}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}-19&-6\\73&23\end{pmatrix}\begin{pmatrix}-41&-12\\147&43\end{pmatrix}$
$14,6,1,1,1,1\ \left(\sqrt{-3}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}3&2\\-11&-7\end{pmatrix}\begin{pmatrix}9&4\\-25&-11\end{pmatrix}\begin{pmatrix}34&25\\-49&-36\end{pmatrix}\begin{pmatrix}19&16\\-25&-21\end{pmatrix}$
$14,6,1,1,1,1\ \left(\sqrt{-3}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}4&3\\-11&-8\end{pmatrix}\begin{pmatrix}14&9\\-25&-16\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}13&4\\-49&-15\end{pmatrix}$
$14,5,2,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-2\\18&7\end{pmatrix}\begin{pmatrix}5&3\\-22&-13\end{pmatrix}\begin{pmatrix}11&9\\-16&-13\end{pmatrix}\begin{pmatrix}39&25\\-64&-41\end{pmatrix}$
$14,5,2,1,1,1\ (\text{cubic})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}-5&-2\\18&7\end{pmatrix}\begin{pmatrix}4&1\\-29&-7\end{pmatrix}\begin{pmatrix}21&4\\-121&-23\end{pmatrix}$
$14,5,2,1,1,1\ (\text{cubic})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}13&4\\-49&-15\end{pmatrix}\begin{pmatrix}4&1\\-29&-7\end{pmatrix}\begin{pmatrix}21&4\\-121&-23\end{pmatrix}$
$14,5,2,1,1,1\ (\text{cubic})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}16&5\\-61&-19\end{pmatrix}\begin{pmatrix}89&25\\-324&-91\end{pmatrix}$
$14,4,3,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}7&4\\-16&-9\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}11&8\\-29&-21\end{pmatrix}\begin{pmatrix}19&16\\-25&-21\end{pmatrix}$
$14,4,2,2,1,1\ \left(\sqrt{-7}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}13&4\\-62&-19\end{pmatrix}\begin{pmatrix}15&4\\-64&-17\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}$
$14,4,2,2,1,1\ \left(\sqrt{-7}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}7&2\\-46&-13\end{pmatrix}\begin{pmatrix}-9&-2\\50&11\end{pmatrix}\begin{pmatrix}29&9\\-100&-31\end{pmatrix}$

$14,3,3,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}11&9\\-16&-13\end{pmatrix}\begin{pmatrix}-3&-1\\22&7\end{pmatrix}\begin{pmatrix}-11&-3\\48&13\end{pmatrix}$
$14,3,3,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}4&3\\-15&-11\end{pmatrix}\begin{pmatrix}-9&-2\\50&11\end{pmatrix}\begin{pmatrix}6&1\\-49&-8\end{pmatrix}$
$14,3,2,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}-5&-1\\46&9\end{pmatrix}\begin{pmatrix}-17&-3\\108&19\end{pmatrix}$
$13,7,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-8&-5\\21&13\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}11&9\\-16&-13\end{pmatrix}\begin{pmatrix}20&9\\-49&-22\end{pmatrix}$
$13,6,2,1,1,1$ $(\sqrt{-3})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}10&3\\-47&-14\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}\begin{pmatrix}32&9\\-121&-34\end{pmatrix}$
$13,6,2,1,1,1$ $(\sqrt{-3})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}7&2\\-39&-11\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}\begin{pmatrix}29&9\\-100&-31\end{pmatrix}$
$13,5,3,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}11&9\\-16&-13\end{pmatrix}\begin{pmatrix}3&1\\-19&-6\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}$
$13,5,2,2,1,1$ $\left(\sqrt{65}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}-19&-8\\50&21\end{pmatrix}\begin{pmatrix}8&3\\-43&-16\end{pmatrix}$
$13,5,2,2,1,1$ $\left(\sqrt{65}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}5&1\\-41&-8\end{pmatrix}\begin{pmatrix}25&4\\-169&-27\end{pmatrix}$
$13,4,3,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}-8&-3\\27&10\end{pmatrix}\begin{pmatrix}2&1\\-17&-8\end{pmatrix}\begin{pmatrix}12&5\\-53&-22\end{pmatrix}$
$13,3,3,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-2&-1\\13&6\end{pmatrix}\begin{pmatrix}-8&-3\\27&10\end{pmatrix}\begin{pmatrix}-7&-6\\13&11\end{pmatrix}\begin{pmatrix}-17&-12\\27&19\end{pmatrix}$
$12,7,2,1,1,1$ $(\sqrt{-3})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}2&1\\-9&-4\end{pmatrix}\begin{pmatrix}-8&-7\\15&13\end{pmatrix}\begin{pmatrix}-24&-17\\41&29\end{pmatrix}\begin{pmatrix}39&25\\-64&-41\end{pmatrix}$
$12,7,2,1,1,1$ $(\sqrt{-3})$	$\begin{pmatrix}-1&-2\\2&3\end{pmatrix}\begin{pmatrix}-1&1\\3&-4\end{pmatrix}\begin{pmatrix}-3&1\\-4&1\end{pmatrix}\begin{pmatrix}-5&16\\-1&3\end{pmatrix}\begin{pmatrix}-11&25\\-4&9\end{pmatrix}$
$12,6,3,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}-8&-3\\27&10\end{pmatrix}\begin{pmatrix}-2&-1\\15&7\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}$
$12,6,2,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-2\\18&7\end{pmatrix}\begin{pmatrix}-11&-8\\18&13\end{pmatrix}\begin{pmatrix}-7&-4\\30&17\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}$
$12,6,2,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&-2\\2&3\end{pmatrix}\begin{pmatrix}3&-2\\2&-1\end{pmatrix}\begin{pmatrix}-1&2\\4&-9\end{pmatrix}\begin{pmatrix}9&4\\-25&-11\end{pmatrix}\begin{pmatrix}-5&16\\-1&3\end{pmatrix}$
$12,5,3,2,1,1$ (quartic)	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}-6&-5\\23&19\end{pmatrix}\begin{pmatrix}-10&-7\\33&23\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}$
$12,5,3,2,1,1$ (quartic)	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}-4&-3\\23&17\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}19&16\\-25&-21\end{pmatrix}$
$12,5,3,2,1,1$ (quartic)	$\begin{pmatrix}2&-1\\-1&1\end{pmatrix}\begin{pmatrix}1&1\\-4&-3\end{pmatrix}\begin{pmatrix}-7&9\\-4&5\end{pmatrix}\begin{pmatrix}6&-13\\1&-2\end{pmatrix}\begin{pmatrix}10&-27\\3&-8\end{pmatrix}$
$12,5,3,2,1,1$ (quartic)	$\begin{pmatrix}2&-1\\-1&1\end{pmatrix}\begin{pmatrix}1&1\\-4&-3\end{pmatrix}\begin{pmatrix}-4&9\\-1&2\end{pmatrix}\begin{pmatrix}11&-13\\6&-7\end{pmatrix}\begin{pmatrix}19&-27\\12&-17\end{pmatrix}$
$12,5,2,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-9&-2\\50&11\end{pmatrix}\begin{pmatrix}-27&-8\\98&29\end{pmatrix}\begin{pmatrix}11&3\\-70&-19\end{pmatrix}$
$12,4,4,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}7&4\\-16&-9\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}-7&-6\\20&17\end{pmatrix}\begin{pmatrix}-17&-12\\44&31\end{pmatrix}$
$12,4,3,3,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}7&5\\-24&-17\end{pmatrix}\begin{pmatrix}-5&-4\\24&19\end{pmatrix}\begin{pmatrix}5&1\\-36&-7\end{pmatrix}$

$12,4,3,3,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -9 & -4 \end{pmatrix} \begin{pmatrix} -8 & -7 \\ 15 & 13 \end{pmatrix} \begin{pmatrix} 23 & 16 \\ -36 & -25 \end{pmatrix} \begin{pmatrix} -44 & -27 \\ 75 & 46 \end{pmatrix}$
$12,4,3,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix} \begin{pmatrix} -11 & -3 \\ 48 & 13 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ -28 & -9 \end{pmatrix} \begin{pmatrix} 17 & 5 \\ -92 & -27 \end{pmatrix}$
$12,4,3,2,2,1$ $\left(\sqrt{-3}\right)$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -9 & 14 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix} \begin{pmatrix} -13 & 16 \\ -9 & 11 \end{pmatrix}$
$12,4,3,2,2,1$ $\left(\sqrt{-3}\right)$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -11 & 14 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix} \begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix}$
$12,3,3,3,2,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix} \begin{pmatrix} 4 & 3 \\ -15 & -11 \end{pmatrix} \begin{pmatrix} -4 & -1 \\ 33 & 8 \end{pmatrix} \begin{pmatrix} -14 & -3 \\ 75 & 16 \end{pmatrix}$
$12,3,3,2,2,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} 7 & -22 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} 13 & -48 \\ 3 & -11 \end{pmatrix}$
$11,9,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 9 \\ -1 & -4 \end{pmatrix} \begin{pmatrix} -7 & -13 \\ 6 & 11 \end{pmatrix} \begin{pmatrix} 34 & 49 \\ -25 & -36 \end{pmatrix} \begin{pmatrix} 39 & 64 \\ -25 & -41 \end{pmatrix}$
$11,8,2,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -2 \\ 18 & 7 \end{pmatrix} \begin{pmatrix} -4 & -3 \\ 19 & 14 \end{pmatrix} \begin{pmatrix} 14 & 9 \\ -25 & -16 \end{pmatrix} \begin{pmatrix} 19 & 16 \\ -25 & -21 \end{pmatrix}$
$11,8,2,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -3 & -2 \\ 17 & 11 \end{pmatrix} \begin{pmatrix} -7 & -4 \\ 23 & 13 \end{pmatrix} \begin{pmatrix} 9 & 4 \\ -25 & -11 \end{pmatrix} \begin{pmatrix} 11 & 9 \\ -16 & -13 \end{pmatrix}$
$11,8,2,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix} \begin{pmatrix} -6 & -5 \\ 17 & 14 \end{pmatrix} \begin{pmatrix} -10 & -7 \\ 23 & 16 \end{pmatrix} \begin{pmatrix} 14 & 9 \\ -25 & -16 \end{pmatrix}$
$11,7,3,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} -2 & -1 \\ 13 & 6 \end{pmatrix} \begin{pmatrix} -13 & -5 \\ 47 & 18 \end{pmatrix} \begin{pmatrix} 29 & 9 \\ -100 & -31 \end{pmatrix}$
$11,7,3,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 17 & 10 \end{pmatrix} \begin{pmatrix} -4 & -3 \\ 19 & 14 \end{pmatrix} \begin{pmatrix} 9 & 4 \\ -25 & -11 \end{pmatrix} \begin{pmatrix} 19 & 16 \\ -25 & -21 \end{pmatrix}$
$11,7,3,1,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -7 & -5 \\ 17 & 12 \end{pmatrix} \begin{pmatrix} -8 & -3 \\ 27 & 10 \end{pmatrix} \begin{pmatrix} 14 & 9 \\ -25 & -16 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ -25 & -6 \end{pmatrix}$
$11,7,2,2,1,1$ $\left(\sqrt{-7}\right)$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 3 & -4 \end{pmatrix} \begin{pmatrix} -3 & 1 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix} \begin{pmatrix} -6 & 25 \\ -1 & 4 \end{pmatrix}$
$11,7,2,2,1,1$ $\left(\sqrt{-7}\right)$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -2 & 1 \\ 5 & -3 \end{pmatrix} \begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} -4 & 1 \\ -9 & 2 \end{pmatrix} \begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix}$
$11,6,4,1,1,1$ $\left(\sqrt{33}\right)$	$\begin{pmatrix} 0 & 1 \\ -1 & -2 \end{pmatrix} \begin{pmatrix} -5 & 4 \\ -4 & 3 \end{pmatrix} \begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} 9 & -2 \\ 5 & -1 \end{pmatrix} \begin{pmatrix} -11 & 4 \\ -25 & 9 \end{pmatrix}$
$11,6,4,1,1,1$ $\left(\sqrt{33}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -9 & -4 \end{pmatrix} \begin{pmatrix} -7 & -6 \\ 13 & 11 \end{pmatrix} \begin{pmatrix} 23 & 16 \\ -36 & -25 \end{pmatrix} \begin{pmatrix} 103 & 64 \\ -169 & -105 \end{pmatrix}$
$11,6,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix} \begin{pmatrix} -11 & -3 \\ 48 & 13 \end{pmatrix} \begin{pmatrix} -3 & -1 \\ 25 & 8 \end{pmatrix} \begin{pmatrix} 21 & 4 \\ -121 & -23 \end{pmatrix}$
$11,6,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -7 & 11 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} -11 & 25 \\ -4 & 9 \end{pmatrix} \begin{pmatrix} -13 & 16 \\ -9 & 11 \end{pmatrix}$
$11,6,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -8 & 11 \\ -3 & 4 \end{pmatrix} \begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} -16 & 25 \\ -9 & 14 \end{pmatrix}$
$11,5,5,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 6 & 11 \\ -5 & -9 \end{pmatrix} \begin{pmatrix} -5 & -18 \\ 2 & 7 \end{pmatrix} \begin{pmatrix} 34 & 49 \\ -25 & -36 \end{pmatrix} \begin{pmatrix} 20 & 49 \\ -9 & -22 \end{pmatrix}$
$11,5,4,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} -2 & -1 \\ 13 & 6 \end{pmatrix} \begin{pmatrix} 18 & 7 \\ -67 & -26 \end{pmatrix} \begin{pmatrix} 14 & 5 \\ -45 & -16 \end{pmatrix}$
$11,5,4,2,1,1$ (cubic)	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} -9 & 14 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix} \begin{pmatrix} -13 & 16 \\ -9 & 11 \end{pmatrix}$

$11,5,4,2,1,1$ (cubic)	$\begin{pmatrix}2&-1\\-1&1\end{pmatrix}\begin{pmatrix}1&1\\-4&-3\end{pmatrix}\begin{pmatrix}-11&14\\-4&5\end{pmatrix}\begin{pmatrix}-9&16\\-4&7\end{pmatrix}\begin{pmatrix}-5&16\\-1&3\end{pmatrix}$
$11,5,3,3,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}4&3\\-15&-11\end{pmatrix}\begin{pmatrix}4&1\\-29&-7\end{pmatrix}\begin{pmatrix}21&4\\-121&-23\end{pmatrix}$
$11,5,3,3,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}-6&-5\\17&14\end{pmatrix}\begin{pmatrix}9&5\\-20&-11\end{pmatrix}\begin{pmatrix}-17&-12\\27&19\end{pmatrix}$
$11,5,3,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&0\\3&-1\end{pmatrix}\begin{pmatrix}-1&-2\\2&3\end{pmatrix}\begin{pmatrix}5&-8\\2&-3\end{pmatrix}\begin{pmatrix}-6&19\\-1&3\end{pmatrix}\begin{pmatrix}-19&81\\-4&17\end{pmatrix}$
$11,4,4,3,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}7&4\\-16&-9\end{pmatrix}\begin{pmatrix}-5&-4\\19&15\end{pmatrix}\begin{pmatrix}11&8\\-29&-21\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}$
$11,4,3,3,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-3&-1\\25&8\end{pmatrix}\begin{pmatrix}15&4\\-64&-17\end{pmatrix}\begin{pmatrix}-14&-3\\75&16\end{pmatrix}$
$10,10,1,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}4&3\\-15&-11\end{pmatrix}\begin{pmatrix}9&4\\-25&-11\end{pmatrix}\begin{pmatrix}14&9\\-25&-16\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}$
$10,10,1,1,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}2&1\\-11&-5\end{pmatrix}\begin{pmatrix}32&9\\-121&-34\end{pmatrix}\begin{pmatrix}29&9\\-100&-31\end{pmatrix}$
$10,10,1,1,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-14&-11\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}14&9\\-25&-16\end{pmatrix}\begin{pmatrix}20&9\\-49&-22\end{pmatrix}$
$10,9,2,1,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}2&1\\-11&-5\end{pmatrix}\begin{pmatrix}-27&-8\\98&29\end{pmatrix}\begin{pmatrix}59&16\\-225&-61\end{pmatrix}$
$10,9,2,1,1,1$ $\left(\sqrt{5}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-3&-1\\22&7\end{pmatrix}\begin{pmatrix}17&4\\-81&-19\end{pmatrix}\begin{pmatrix}32&9\\-121&-34\end{pmatrix}$
$10,8,3,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-14&-11\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}-13&-8\\31&19\end{pmatrix}\begin{pmatrix}27&16\\-49&-29\end{pmatrix}$
$10,7,4,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}4&3\\-15&-11\end{pmatrix}\begin{pmatrix}-8&-5\\21&13\end{pmatrix}\begin{pmatrix}4&1\\-25&-6\end{pmatrix}\begin{pmatrix}20&9\\-49&-22\end{pmatrix}$
$10,7,3,2,1,1$ $\left(\sqrt{21}\right)$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-9&-7\end{pmatrix}\begin{pmatrix}2&1\\-11&-5\end{pmatrix}\begin{pmatrix}-19&-6\\73&23\end{pmatrix}\begin{pmatrix}-41&-12\\147&43\end{pmatrix}$
$10,7,3,2,1,1$ $\left(\sqrt{21}\right)$	$\begin{pmatrix}-3&2\\1&-1\end{pmatrix}\begin{pmatrix}-2&-3\\3&4\end{pmatrix}\begin{pmatrix}5&-8\\2&-3\end{pmatrix}\begin{pmatrix}2&1\\-9&-4\end{pmatrix}\begin{pmatrix}-5&16\\-1&3\end{pmatrix}$
$10,7,2,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-2\\18&7\end{pmatrix}\begin{pmatrix}-11&-8\\18&13\end{pmatrix}\begin{pmatrix}-5&-3\\32&19\end{pmatrix}\begin{pmatrix}-9&-5\\38&21\end{pmatrix}$
$10,6,5,1,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-14&-11\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}9&5\\-20&-11\end{pmatrix}\begin{pmatrix}39&25\\-64&-41\end{pmatrix}$
$10,6,5,1,1,1$ (cubic)	$\begin{pmatrix}1&0\\-1&1\end{pmatrix}\begin{pmatrix}2&9\\-1&-4\end{pmatrix}\begin{pmatrix}6&11\\-5&-9\end{pmatrix}\begin{pmatrix}42&61\\-31&-45\end{pmatrix}\begin{pmatrix}233&324\\-169&-235\end{pmatrix}$
$10,6,5,1,1,1$ (cubic)	$\begin{pmatrix}2&-1\\-1&1\end{pmatrix}\begin{pmatrix}1&1\\-4&-3\end{pmatrix}\begin{pmatrix}-8&11\\-3&4\end{pmatrix}\begin{pmatrix}-5&16\\-1&3\end{pmatrix}\begin{pmatrix}-16&25\\-9&14\end{pmatrix}$
$10,6,5,1,1,1$ (cubic)	$\begin{pmatrix}2&-1\\-1&1\end{pmatrix}\begin{pmatrix}1&1\\-4&-3\end{pmatrix}\begin{pmatrix}-7&11\\-2&3\end{pmatrix}\begin{pmatrix}-11&25\\-4&9\end{pmatrix}\begin{pmatrix}-13&16\\-9&11\end{pmatrix}$
$10,6,4,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}5&4\\-14&-11\end{pmatrix}\begin{pmatrix}3&1\\-16&-5\end{pmatrix}\begin{pmatrix}-19&-12\\46&29\end{pmatrix}\begin{pmatrix}-11&-6\\24&13\end{pmatrix}$
$10,6,4,2,1,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-3&-2\\8&5\end{pmatrix}\begin{pmatrix}-3&-1\\22&7\end{pmatrix}\begin{pmatrix}15&4\\-64&-17\end{pmatrix}\begin{pmatrix}41&9\\-196&-43\end{pmatrix}$
$10,6,3,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix}-1&1\\0&-1\end{pmatrix}\begin{pmatrix}-5&-3\\12&7\end{pmatrix}\begin{pmatrix}-7&-2\\32&9\end{pmatrix}\begin{pmatrix}-23&-18\\32&25\end{pmatrix}\begin{pmatrix}-7&-5\\38&27\end{pmatrix}$

$10,5,5,2,1,1\ \left(\sqrt{5}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 4 & 3 \\ -15 & -11 \end{pmatrix}\begin{pmatrix} 11 & 7 \\ -30 & -19 \end{pmatrix}\begin{pmatrix} 9 & 5 \\ -20 & -11 \end{pmatrix}\begin{pmatrix} 4 & 1 \\ -25 & -6 \end{pmatrix}$
$10,5,5,2,1,1\ \left(\sqrt{5}\right)$	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}\begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix}\begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix}\begin{pmatrix} -6 & 19 \\ -1 & 3 \end{pmatrix}\begin{pmatrix} -19 & 81 \\ -4 & 17 \end{pmatrix}$
$10,5,4,2,2,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix}\begin{pmatrix} -3 & -1 \\ 22 & 7 \end{pmatrix}\begin{pmatrix} 25 & 7 \\ -118 & -33 \end{pmatrix}\begin{pmatrix} 19 & 5 \\ -80 & -21 \end{pmatrix}$
$10,5,4,2,2,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix}\begin{pmatrix} -10 & 7 \\ -3 & 2 \end{pmatrix}\begin{pmatrix} -16 & 9 \\ -25 & 14 \end{pmatrix}$
$10,5,4,2,2,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix}\begin{pmatrix} -12 & 7 \\ -7 & 4 \end{pmatrix}\begin{pmatrix} -6 & 5 \\ -5 & 4 \end{pmatrix}$
$10,5,3,2,2,2\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix}\begin{pmatrix} 13 & -18 \\ 8 & -11 \end{pmatrix}\begin{pmatrix} -13 & 22 \\ -3 & 5 \end{pmatrix}$
$10,4,4,3,2,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix}\begin{pmatrix} 7 & 5 \\ -24 & -17 \end{pmatrix}\begin{pmatrix} -1 & -1 \\ 10 & 9 \end{pmatrix}\begin{pmatrix} -9 & -7 \\ 40 & 31 \end{pmatrix}$
$10,4,3,3,2,2\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 7 & -12 \\ 3 & -5 \end{pmatrix}\begin{pmatrix} -7 & 8 \\ -1 & 1 \end{pmatrix}\begin{pmatrix} -17 & 24 \\ -5 & 7 \end{pmatrix}$
$9,9,2,2,1,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -5 & -2 \\ 18 & 7 \end{pmatrix}\begin{pmatrix} -11 & -8 \\ 18 & 13 \end{pmatrix}\begin{pmatrix} 5 & 3 \\ -27 & -16 \end{pmatrix}\begin{pmatrix} 17 & 4 \\ -81 & -19 \end{pmatrix}$
$9,9,2,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -19 & -6 \end{pmatrix}\begin{pmatrix} -35 & -8 \\ 162 & 37 \end{pmatrix}\begin{pmatrix} 75 & 16 \\ -361 & -77 \end{pmatrix}$
$9,9,2,2,1,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix}\begin{pmatrix} 2 & -5 \\ -5 & 13 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix}\begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix}$
$9,9,2,2,1,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} -3 & 1 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} 1 & -1 \\ -5 & 6 \end{pmatrix}\begin{pmatrix} 3 & -5 \\ -7 & 12 \end{pmatrix}\begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix}$
$9,8,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 3 & 2 \\ -11 & -7 \end{pmatrix}\begin{pmatrix} 11 & 9 \\ -16 & -13 \end{pmatrix}\begin{pmatrix} -13 & -6 \\ 37 & 17 \end{pmatrix}\begin{pmatrix} -29 & -12 \\ 75 & 31 \end{pmatrix}$
$9,8,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} -3 & 1 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} -1 & 1 \\ 4 & -5 \end{pmatrix}\begin{pmatrix} 2 & -5 \\ -5 & 13 \end{pmatrix}\begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix}$
$9,8,3,2,1,1$ (cubic)	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 4 & -3 \\ 3 & -2 \end{pmatrix}\begin{pmatrix} -3 & 1 \\ 8 & -3 \end{pmatrix}\begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix}$
$9,7,5,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} 3 & 2 \\ -11 & -7 \end{pmatrix}\begin{pmatrix} 11 & 9 \\ -16 & -13 \end{pmatrix}\begin{pmatrix} 11 & 5 \\ -31 & -14 \end{pmatrix}\begin{pmatrix} 64 & 25 \\ -169 & -66 \end{pmatrix}$
$9,7,4,2,1,1\ \left(\sqrt{-7}\right)$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} 7 & -5 \\ 3 & -2 \end{pmatrix}\begin{pmatrix} -7 & 9 \\ -4 & 5 \end{pmatrix}\begin{pmatrix} -5 & 16 \\ -1 & 3 \end{pmatrix}$
$9,7,4,2,1,1\ \left(\sqrt{-7}\right)$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix}\begin{pmatrix} 8 & -5 \\ 5 & -3 \end{pmatrix}\begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix}\begin{pmatrix} -13 & 16 \\ -9 & 11 \end{pmatrix}$
$9,7,3,2,2,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -19 & -6 \end{pmatrix}\begin{pmatrix} -25 & -6 \\ 121 & 29 \end{pmatrix}\begin{pmatrix} -53 & -12 \\ 243 & 55 \end{pmatrix}$
$9,7,3,2,2,1$ (cubic)	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 6 & -7 \\ 1 & -1 \end{pmatrix}\begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix}\begin{pmatrix} -16 & 25 \\ -9 & 14 \end{pmatrix}$
$9,7,3,2,2,1$ (cubic)	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}\begin{pmatrix} 6 & -7 \\ 1 & -1 \end{pmatrix}\begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix}\begin{pmatrix} -16 & 25 \\ -9 & 14 \end{pmatrix}$
$9,6,5,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\begin{pmatrix} -3 & -2 \\ 8 & 5 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -19 & -6 \end{pmatrix}\begin{pmatrix} 21 & 5 \\ -101 & -24 \end{pmatrix}\begin{pmatrix} 114 & 25 \\ -529 & -116 \end{pmatrix}$
$9,6,5,2,1,1$ (cubic)	$\begin{pmatrix} -1 & 2 \\ 1 & -3 \end{pmatrix}\begin{pmatrix} 1 & 4 \\ -1 & -3 \end{pmatrix}\begin{pmatrix} 5 & -2 \\ 8 & -3 \end{pmatrix}\begin{pmatrix} 4 & -3 \\ -9 & 7 \end{pmatrix}\begin{pmatrix} 3 & 1 \\ -16 & -5 \end{pmatrix}$

9, 6, 5, 2, 1, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}-3 & 4\\ -1 & 1\end{pmatrix}\begin{pmatrix}1 & 0\\ -5 & 1\end{pmatrix}\begin{pmatrix}4 & -3\\ -9 & 7\end{pmatrix}\begin{pmatrix}-16 & 9\\ -25 & 14\end{pmatrix}$
9, 6, 4, 3, 1, 1 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}-5 & -3\\ 12 & 7\end{pmatrix}\begin{pmatrix}7 & 5\\ -24 & -17\end{pmatrix}\begin{pmatrix}1 & 1\\ -9 & -8\end{pmatrix}\begin{pmatrix}17 & 4\\ -81 & -19\end{pmatrix}$
9, 6, 3, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & 0\\ 3 & -1\end{pmatrix}\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}7 & -12\\ 3 & -5\end{pmatrix}\begin{pmatrix}6 & -7\\ 1 & -1\end{pmatrix}\begin{pmatrix}-15 & 49\\ -4 & 13\end{pmatrix}$
9, 6, 3, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}4 & -3\\ 3 & -2\end{pmatrix}\begin{pmatrix}-8 & 3\\ -3 & 1\end{pmatrix}\begin{pmatrix}1 & -3\\ -3 & 10\end{pmatrix}\begin{pmatrix}9 & 4\\ -25 & -11\end{pmatrix}$
9, 6, 3, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}-3 & 4\\ -1 & 1\end{pmatrix}\begin{pmatrix}-1 & 0\\ 6 & -1\end{pmatrix}\begin{pmatrix}4 & -3\\ -9 & 7\end{pmatrix}\begin{pmatrix}7 & -3\\ 12 & -5\end{pmatrix}$
9, 5, 5, 2, 2, 1 ($\mathbb{Q}$)	$\begin{pmatrix}2 & -1\\ -1 & 1\end{pmatrix}\begin{pmatrix}1 & 1\\ -4 & -3\end{pmatrix}\begin{pmatrix}7 & -18\\ 2 & -5\end{pmatrix}\begin{pmatrix}13 & -18\\ 8 & -11\end{pmatrix}\begin{pmatrix}-13 & 22\\ -3 & 5\end{pmatrix}$
9, 5, 4, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}1 & 0\\ -4 & 1\end{pmatrix}\begin{pmatrix}4 & -3\\ 3 & -2\end{pmatrix}\begin{pmatrix}-5 & 11\\ -1 & 2\end{pmatrix}\begin{pmatrix}-15 & 49\\ -4 & 13\end{pmatrix}$
9, 5, 4, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}4 & -3\\ 3 & -2\end{pmatrix}\begin{pmatrix}1 & 0\\ -5 & 1\end{pmatrix}\begin{pmatrix}2 & -5\\ -5 & 13\end{pmatrix}\begin{pmatrix}-5 & 16\\ -1 & 3\end{pmatrix}$
9, 5, 4, 3, 2, 1 (cubic)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}1 & 0\\ -5 & 1\end{pmatrix}\begin{pmatrix}-5 & 4\\ -4 & 3\end{pmatrix}\begin{pmatrix}3 & -5\\ -7 & 12\end{pmatrix}\begin{pmatrix}-4 & 9\\ -1 & 2\end{pmatrix}$
9, 5, 3, 3, 3, 1 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}-5 & -3\\ 12 & 7\end{pmatrix}\begin{pmatrix}1 & 1\\ -9 & -8\end{pmatrix}\begin{pmatrix}-10 & -7\\ 33 & 23\end{pmatrix}\begin{pmatrix}-11 & -3\\ 48 & 13\end{pmatrix}$
9, 4, 3, 3, 3, 2 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & 0\\ 3 & -1\end{pmatrix}\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}6 & -7\\ 1 & -1\end{pmatrix}\begin{pmatrix}-9 & 16\\ -4 & 7\end{pmatrix}\begin{pmatrix}10 & -27\\ 3 & -8\end{pmatrix}$
8, 8, 4, 2, 1, 1 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}7 & 4\\ -16 & -9\end{pmatrix}\begin{pmatrix}1 & 1\\ -8 & -7\end{pmatrix}\begin{pmatrix}7 & 5\\ -24 & -17\end{pmatrix}\begin{pmatrix}23 & 9\\ -64 & -25\end{pmatrix}$
8, 8, 3, 3, 1, 1 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}-5 & -3\\ 12 & 7\end{pmatrix}\begin{pmatrix}1 & 1\\ -8 & -7\end{pmatrix}\begin{pmatrix}11 & 8\\ -40 & -29\end{pmatrix}\begin{pmatrix}29 & 9\\ -100 & -31\end{pmatrix}$
8, 8, 3, 3, 1, 1 ($\sqrt{-2}$)	$\begin{pmatrix}-1 & 0\\ 3 & -1\end{pmatrix}\begin{pmatrix}-2 & -3\\ 3 & 4\end{pmatrix}\begin{pmatrix}3 & -8\\ -1 & 3\end{pmatrix}\begin{pmatrix}-7 & 9\\ -4 & 5\end{pmatrix}\begin{pmatrix}3 & 16\\ -1 & -5\end{pmatrix}$
8, 8, 3, 3, 1, 1 ($\sqrt{-2}$)	$\begin{pmatrix}-1 & 0\\ 3 & -1\end{pmatrix}\begin{pmatrix}-2 & -3\\ 3 & 4\end{pmatrix}\begin{pmatrix}4 & -7\\ -1 & 2\end{pmatrix}\begin{pmatrix}-4 & 9\\ -1 & 2\end{pmatrix}\begin{pmatrix}9 & 25\\ -4 & -11\end{pmatrix}$
8, 8, 2, 2, 2, 2 ($\mathbb{Q}$)	$\begin{pmatrix}-1 & -2\\ 2 & 3\end{pmatrix}\begin{pmatrix}3 & -2\\ 2 & -1\end{pmatrix}\begin{pmatrix}-5 & -2\\ 18 & 7\end{pmatrix}\begin{pmatrix}7 & -18\\ 2 & -5\end{pmatrix}\begin{pmatrix}1 & -4\\ -4 & 17\end{pmatrix}$
8, 7, 6, 1, 1, 1 ($\sqrt{-3}$)	$\begin{pmatrix}3 & -1\\ -2 & 1\end{pmatrix}\begin{pmatrix}1 & 1\\ -4 & -3\end{pmatrix}\begin{pmatrix}3 & -8\\ -1 & 3\end{pmatrix}\begin{pmatrix}-7 & 9\\ -4 & 5\end{pmatrix}\begin{pmatrix}3 & 16\\ -1 & -5\end{pmatrix}$
8, 7, 6, 1, 1, 1 ($\sqrt{-3}$)	$\begin{pmatrix}3 & -1\\ -2 & 1\end{pmatrix}\begin{pmatrix}1 & 1\\ -4 & -3\end{pmatrix}\begin{pmatrix}4 & -7\\ -1 & 2\end{pmatrix}\begin{pmatrix}-4 & 9\\ -1 & 2\end{pmatrix}\begin{pmatrix}9 & 25\\ -4 & -11\end{pmatrix}$
8, 7, 5, 2, 1, 1 (cubic)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}-8 & -5\\ 21 & 13\end{pmatrix}\begin{pmatrix}1 & 1\\ -8 & -7\end{pmatrix}\begin{pmatrix}7 & 5\\ -24 & -17\end{pmatrix}\begin{pmatrix}20 & 9\\ -49 & -22\end{pmatrix}$
8, 7, 5, 2, 1, 1 (cubic)	$\begin{pmatrix}2 & -1\\ -1 & 1\end{pmatrix}\begin{pmatrix}1 & 1\\ -4 & -3\end{pmatrix}\begin{pmatrix}6 & -7\\ 1 & -1\end{pmatrix}\begin{pmatrix}10 & -17\\ 3 & -5\end{pmatrix}\begin{pmatrix}-11 & 25\\ -4 & 9\end{pmatrix}$
8, 7, 5, 2, 1, 1 (cubic)	$\begin{pmatrix}2 & -1\\ -1 & 1\end{pmatrix}\begin{pmatrix}1 & 1\\ -4 & -3\end{pmatrix}\begin{pmatrix}6 & -7\\ 1 & -1\end{pmatrix}\begin{pmatrix}7 & -18\\ 2 & -5\end{pmatrix}\begin{pmatrix}-16 & 25\\ -9 & 14\end{pmatrix}$
8, 7, 4, 3, 1, 1 ($\sqrt{-6}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}7 & 4\\ -16 & -9\end{pmatrix}\begin{pmatrix}1 & 1\\ -8 & -7\end{pmatrix}\begin{pmatrix}11 & 8\\ -29 & -21\end{pmatrix}\begin{pmatrix}13 & 4\\ -49 & -15\end{pmatrix}$
8, 7, 4, 3, 1, 1 ($\sqrt{-6}$)	$\begin{pmatrix}-1 & 1\\ 0 & -1\end{pmatrix}\begin{pmatrix}7 & 4\\ -16 & -9\end{pmatrix}\begin{pmatrix}1 & 1\\ -8 & -7\end{pmatrix}\begin{pmatrix}-8 & -3\\ 27 & 10\end{pmatrix}\begin{pmatrix}34 & 25\\ -49 & -36\end{pmatrix}$

$8,7,4,2,2,1\ (\sqrt{-7})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -3 & 4 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ -7 & 5 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 7 & -1 \end{pmatrix} \begin{pmatrix} -5 & 2 \\ 17 & -7 \end{pmatrix}$
$8,7,4,2,2,1\ (\sqrt{-7})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & -2 \\ -3 & 7 \end{pmatrix} \begin{pmatrix} -7 & 2 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -5 & 4 \\ -4 & 3 \end{pmatrix} \begin{pmatrix} 9 & 4 \\ -25 & -11 \end{pmatrix}$
$8,7,3,3,2,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 5 & -6 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} 13 & -21 \\ 5 & -8 \end{pmatrix} \begin{pmatrix} -22 & 49 \\ -9 & 20 \end{pmatrix}$
$8,6,6,2,1,1\ (\sqrt{-3})$	$\begin{pmatrix} -2 & 3 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} -4 & 1 \\ 7 & -2 \end{pmatrix} \begin{pmatrix} -4 & 3 \\ 5 & -4 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -9 & -4 \end{pmatrix} \begin{pmatrix} 2 & 9 \\ -1 & -4 \end{pmatrix}$
$8,6,6,2,1,1\ (\sqrt{-3})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 6 & 5 \end{pmatrix} \begin{pmatrix} -17 & -10 \\ 46 & 27 \end{pmatrix} \begin{pmatrix} -19 & -8 \\ 50 & 21 \end{pmatrix} \begin{pmatrix} 39 & 25 \\ -64 & -41 \end{pmatrix}$
$8,6,5,3,1,1\ \left(\sqrt{5}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} -10 & -7 \\ 33 & 23 \end{pmatrix} \begin{pmatrix} 32 & 9 \\ -121 & -34 \end{pmatrix}$
$8,6,5,3,1,1\ \left(\sqrt{5}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} -10 & -7 \\ 33 & 23 \end{pmatrix} \begin{pmatrix} 32 & 9 \\ -121 & -34 \end{pmatrix}$
$8,6,5,2,2,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} 7 & 5 \\ -24 & -17 \end{pmatrix} \begin{pmatrix} 11 & 7 \\ -30 & -19 \end{pmatrix} \begin{pmatrix} 9 & 5 \\ -20 & -11 \end{pmatrix}$
$8,6,4,3,2,1\ \left(\sqrt{2}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} -15 & -11 \\ 56 & 41 \end{pmatrix} \begin{pmatrix} -13 & -9 \\ 42 & 29 \end{pmatrix}$
$8,6,4,3,2,1\ \left(\sqrt{2}\right)$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 5 & -6 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix} \begin{pmatrix} -25 & 64 \\ -9 & 23 \end{pmatrix}$
$8,6,4,2,2,2\ (\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} 11 & -6 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 13 & -10 \\ 4 & -3 \end{pmatrix}$
$8,5,4,3,3,1\ \left(\sqrt{10}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} -8 & -3 \\ 27 & 10 \end{pmatrix} \begin{pmatrix} 9 & 5 \\ -20 & -11 \end{pmatrix} \begin{pmatrix} -17 & -12 \\ 27 & 19 \end{pmatrix}$
$8,5,4,3,3,1\ \left(\sqrt{10}\right)$	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} 6 & -7 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix} \begin{pmatrix} 10 & -27 \\ 3 & -8 \end{pmatrix}$
$8,5,4,3,2,2\ (\mathbb{Q})$	$\begin{pmatrix} 1 & -1 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 5 & -1 \\ 6 & -1 \end{pmatrix} \begin{pmatrix} -19 & 7 \\ -30 & 11 \end{pmatrix} \begin{pmatrix} -11 & 5 \\ -20 & 9 \end{pmatrix}$
$8,4,4,4,3,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 7 & 4 \\ -16 & -9 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -8 & -7 \end{pmatrix} \begin{pmatrix} 11 & 8 \\ -40 & -29 \end{pmatrix} \begin{pmatrix} 11 & 4 \\ -36 & -13 \end{pmatrix}$
$8,4,4,4,2,2\ (\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -5 & 4 \\ -4 & 3 \end{pmatrix} \begin{pmatrix} -9 & 14 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 7 & -18 \\ 2 & -5 \end{pmatrix}$
$7,7,7,1,1,1\ (\mathbb{Q})$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 7 & 6 \end{pmatrix} \begin{pmatrix} -8 & -5 \\ 21 & 13 \end{pmatrix} \begin{pmatrix} 13 & 4 \\ -49 & -15 \end{pmatrix} \begin{pmatrix} 20 & 9 \\ -49 & -22 \end{pmatrix}$
$7,7,5,3,1,1\ \left(\sqrt{21}\right)$	$\begin{pmatrix} 1 & -3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} -2 & -3 \\ 5 & 7 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ -9 & -7 \end{pmatrix} \begin{pmatrix} -7 & 4 \\ -9 & 5 \end{pmatrix}$
$7,7,5,3,1,1\ \left(\sqrt{21}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -5 & -3 \\ 12 & 7 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 7 & 6 \end{pmatrix} \begin{pmatrix} -27 & -8 \\ 98 & 29 \end{pmatrix} \begin{pmatrix} 59 & 16 \\ -225 & -61 \end{pmatrix}$
$7,7,4,4,1,1\ (\sqrt{-7})$	$\begin{pmatrix} -2 & 1 \\ 5 & -3 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ -3 & 5 \end{pmatrix} \begin{pmatrix} 1 & -4 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 5 & 9 \\ -4 & -7 \end{pmatrix} \begin{pmatrix} -4 & 1 \\ -9 & 2 \end{pmatrix}$
$7,7,4,4,1,1\ (\sqrt{-7})$	$\begin{pmatrix} -2 & 1 \\ 5 & -3 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ -3 & 5 \end{pmatrix} \begin{pmatrix} 1 & -4 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 5 & 9 \\ -4 & -7 \end{pmatrix} \begin{pmatrix} -4 & 1 \\ -9 & 2 \end{pmatrix}$
$7,7,3,3,2,2\ \left(\sqrt{7}\right)$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} -1 & 2 \\ 5 & -11 \end{pmatrix} \begin{pmatrix} 1 & -5 \\ -2 & 11 \end{pmatrix} \begin{pmatrix} -8 & -3 \\ 27 & 10 \end{pmatrix}$
$7,7,3,3,2,2\ \left(\sqrt{7}\right)$	$\begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -4 & 5 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 17 & -37 \\ 6 & -13 \end{pmatrix} \begin{pmatrix} 31 & -75 \\ 12 & -29 \end{pmatrix}$

$7,6,6,2,2,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} -5 & 3 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 6 & -1 \end{pmatrix} \begin{pmatrix} -5 & 2 \\ 12 & -5 \end{pmatrix} \begin{pmatrix} -7 & 9 \\ -4 & 5 \end{pmatrix}$
$7,6,5,4,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} 5 & -6 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix} \begin{pmatrix} -25 & 64 \\ -9 & 23 \end{pmatrix}$
$7,6,5,3,2,1$ (cubic)	$\begin{pmatrix} 0 & 1 \\ -1 & -3 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix} \begin{pmatrix} -4 & 5 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} -14 & 31 \\ -5 & 11 \end{pmatrix} \begin{pmatrix} -66 & 169 \\ -25 & 64 \end{pmatrix}$
$7,6,5,3,2,1$ (cubic)	$\begin{pmatrix} 1 & -3 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -3 & 1 \\ 5 & -2 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ -9 & -4 \end{pmatrix} \begin{pmatrix} -9 & 5 \\ 7 & -4 \end{pmatrix} \begin{pmatrix} -9 & 7 \\ 5 & -4 \end{pmatrix}$
$7,6,5,3,2,1$ (cubic)	$\begin{pmatrix} -2 & -3 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} -3 & 5 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} 3 & -2 \\ -7 & 5 \end{pmatrix} \begin{pmatrix} -4 & 9 \\ -1 & 2 \end{pmatrix}$
$7,6,4,4,2,1$ $(\mathbb{Q})$	$\begin{pmatrix} 3 & -1 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} -5 & 7 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} 7 & -17 \\ -2 & 5 \end{pmatrix} \begin{pmatrix} 1 & -7 \\ 0 & 1 \end{pmatrix}$
$7,5,5,4,2,1$ $\left(\sqrt{2}\right)$	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} 5 & -6 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} -19 & 30 \\ -7 & 11 \end{pmatrix} \begin{pmatrix} -11 & 20 \\ -5 & 9 \end{pmatrix}$
$7,5,5,4,2,1$ $\left(\sqrt{2}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 7 & 6 \end{pmatrix} \begin{pmatrix} 7 & 4 \\ -16 & -9 \end{pmatrix} \begin{pmatrix} 18 & 7 \\ -67 & -26 \end{pmatrix} \begin{pmatrix} 14 & 5 \\ -45 & -16 \end{pmatrix}$
$7,5,5,3,2,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 4 & -3 \\ 3 & -2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} 3 & -7 \\ -8 & 19 \end{pmatrix} \begin{pmatrix} 1 & -5 \\ -2 & 11 \end{pmatrix}$
$7,5,4,4,3,1$ $(\mathbb{Q})$	$\begin{pmatrix} 0 & 1 \\ -1 & -2 \end{pmatrix} \begin{pmatrix} 5 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -6 & 5 \\ -5 & 4 \end{pmatrix} \begin{pmatrix} -19 & 11 \\ -7 & 4 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix}$
$7,5,4,3,3,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} 9 & -5 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 7 & -12 \\ 3 & -5 \end{pmatrix} \begin{pmatrix} -6 & 5 \\ -5 & 4 \end{pmatrix}$
$6,6,6,4,1,1$ $(\mathbb{Q})$	$\begin{pmatrix} 3 & -1 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} -5 & 7 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} -1 & 6 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -11 & 25 \\ -4 & 9 \end{pmatrix}$
$6,6,6,2,2,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 5 & -8 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ 8 & -3 \end{pmatrix} \begin{pmatrix} -3 & 2 \\ 10 & -7 \end{pmatrix} \begin{pmatrix} -1 & 4 \\ 2 & -9 \end{pmatrix}$
$6,6,5,5,1,1$ $\left(\sqrt{3}\right)$	$\begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} -4 & 5 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} -14 & 31 \\ -5 & 11 \end{pmatrix} \begin{pmatrix} -66 & 169 \\ -25 & 64 \end{pmatrix}$
$6,6,5,5,1,1$ $\left(\sqrt{3}\right)$	$\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} -1 & -1 \\ 6 & 5 \end{pmatrix} \begin{pmatrix} 9 & 5 \\ -20 & -11 \end{pmatrix} \begin{pmatrix} 14 & 9 \\ -39 & -25 \end{pmatrix} \begin{pmatrix} 64 & 25 \\ -169 & -66 \end{pmatrix}$
$6,6,5,4,2,1$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -7 & 4 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} -8 & 11 \\ -3 & 4 \end{pmatrix} \begin{pmatrix} -16 & 25 \\ -9 & 14 \end{pmatrix}$
$6,6,5,3,3,1$ $(\mathbb{Q})$	$\begin{pmatrix} 3 & -1 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -4 & -3 \end{pmatrix} \begin{pmatrix} 7 & -9 \\ -3 & 4 \end{pmatrix} \begin{pmatrix} -1 & 6 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 7 & -12 \\ 3 & -5 \end{pmatrix}$
$6,6,4,4,2,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} -7 & 4 \\ -2 & 1 \end{pmatrix} \begin{pmatrix} -11 & 14 \\ -4 & 5 \end{pmatrix} \begin{pmatrix} -9 & 16 \\ -4 & 7 \end{pmatrix}$
$6,6,4,3,3,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 4 & -3 \\ 3 & -2 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 6 & -1 \end{pmatrix} \begin{pmatrix} -1 & 4 \\ 2 & -9 \end{pmatrix} \begin{pmatrix} -12 & 5 \\ -5 & 2 \end{pmatrix}$
$6,6,3,3,3,3$ $(\mathbb{Q})$	$\begin{pmatrix} -2 & -3 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 4 & -3 \\ 3 & -2 \end{pmatrix} \begin{pmatrix} 1 & 3 \\ -3 & -8 \end{pmatrix} \begin{pmatrix} -8 & 3 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 1 & -3 \\ -3 & 10 \end{pmatrix}$
$6,5,4,4,3,2$ $(\mathbb{Q})$	$\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} -5 & 4 \\ -4 & 3 \end{pmatrix} \begin{pmatrix} -1 & 4 \\ 2 & -9 \end{pmatrix} \begin{pmatrix} 7 & -12 \\ 3 & -5 \end{pmatrix}$
$5,5,5,5,2,2$ $(\mathbb{Q})$	$\begin{pmatrix} 3 & -2 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & -2 \\ -3 & 7 \end{pmatrix} \begin{pmatrix} 1 & 3 \\ -3 & -8 \end{pmatrix} \begin{pmatrix} 4 & 7 \\ -7 & -12 \end{pmatrix} \begin{pmatrix} 4 & 5 \\ -5 & -6 \end{pmatrix}$
$5,5,4,4,3,3$ $(\mathbb{Q})$	$\begin{pmatrix} 1 & 0 \\ -4 & 1 \end{pmatrix} \begin{pmatrix} 4 & -3 \\ 3 & -2 \end{pmatrix} \begin{pmatrix} 3 & 4 \\ -4 & -5 \end{pmatrix} \begin{pmatrix} -1 & 5 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 3 & 8 \\ -5 & -13 \end{pmatrix}$

$4,4,4,4,4,4$ $(\mathbb{Q})$	$\begin{pmatrix}1&0\\-4&1\end{pmatrix}\begin{pmatrix}1&-4\\0&1\end{pmatrix}\begin{pmatrix}3&4\\-4&-5\end{pmatrix}\begin{pmatrix}-5&4\\-4&3\end{pmatrix}\begin{pmatrix}5&-8\\-8&13\end{pmatrix}$

4 Conclusions and Outlook

In this paper, we have further developed the connections between the torsion-free, genus zero modular subgroups and the theory of dessins d'enfants. This was accomplished through construction of the ramification data and the dessins for each subgroup. In addition, we have shown that the index 24 and 36 subgroups are of special interest, since they are naturally associated to K3 surfaces and Calabi-Yau threefolds, respectively.

In a parallel vein, we have found the modular subgroups associated with all the 112 semi-stable elliptic fibrations over $\mathbb{P}^1$ which are extremal K3 surfaces with six singular fibres. Indeed, at the intersection of the 112 and 33 lie the nine K3 surfaces. This result is important not only for developing our understanding of these surfaces and their connection to the theory of modular subgroups, but also because these connections should give us further tools for categorizing these surfaces, and the relations between them.

To our intricate web of correspondences we can add physics. In [18], Seiberg-Witten curves for four-dimensional supersymmetric field theories with unitary gauge groups were given an interpretation as clean dessins. In [3], the nine K3 surfaces were singled out to relate to the recently advanced class of $\mathcal{N} = 2$ Gaiotto theories. Now, we have an extensive catalogue of dessins which have underlying Calabi-Yau structure, to construct their associated gauge theory is clearly an interesting story which we are currently pursuing [19].

A Dessins from the Miranda-Persson Table

In this appendix we give the dessins for all the entires in [10]. These dessins match those already given in [11], hence their being relegated to an appendix. Nevertheless, there are two advantages to our drawings over those in [11]: first, they are drawn electronically; second, we have marked the black/white nodes as an emphasis of the bipartite nature of the graphs.



$19, 1, 1, 1, 1, 1\ (\mathbb{Q})$



$18, 2, 1, 1, 1, 1\ (\mathbb{Q})$



$18, 2, 1, 1, 1, 1\ (\sqrt{-3})$



$18, 2, 1, 1, 1, 1\ (\sqrt{-3})$



$17, 3, 1, 1, 1, 1\ (\mathbb{Q})$



$17, 2, 2, 1, 1, 1\ \left(\sqrt{17}\right)$



$17, 2, 2, 1, 1, 1\ \left(\sqrt{17}\right)$



$16, 4, 1, 1, 1, 1\ (\mathbb{Q})$



$16, 3, 2, 1, 1, 1\ (\mathbb{Q})$



$16, 3, 2, 1, 1, 1\ (\sqrt{-2})$



$16, 3, 2, 1, 1, 1\ (\sqrt{-2})$



$16, 2, 2, 2, 1, 1\ (\mathbb{Q})$



$15, 5, 1, 1, 1, 1$ $(\sqrt{-15})$



$15, 5, 1, 1, 1, 1$ $(\sqrt{-15})$



$15, 4, 2, 1, 1, 1$ $(\sqrt{-15})$



$15, 4, 2, 1, 1, 1$ $(\sqrt{-15})$



$15, 3, 3, 1, 1, 1$ $(\mathbb{Q})$



$15, 3, 2, 2, 1, 1$ $(\mathbb{Q})$



$15, 3, 2, 2, 1, 1$ $(\sqrt{-15})$



$15, 3, 2, 2, 1, 1$ $(\sqrt{-15})$



$14, 6, 1, 1, 1, 1$ $(\sqrt{-3})$



$14, 6, 1, 1, 1, 1$ $(\sqrt{-3})$



$14, 5, 2, 1, 1, 1$ $(\mathbb{Q})$



$14, 5, 2, 1, 1, 1$ (cubic)



$14, 5, 2, 1, 1, 1$ (cubic)



$14, 5, 2, 1, 1, 1$ (cubic)



$14, 4, 3, 1, 1, 1$ ($\mathbb{Q}$)



$14, 4, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$



$14, 4, 2, 2, 1, 1$ $\left(\sqrt{-7}\right)$



$14, 3, 3, 2, 1, 1$ ($\mathbb{Q}$)



$14, 3, 3, 2, 1, 1$ ($\mathbb{Q}$)



$14, 3, 2, 2, 2, 1$ ($\mathbb{Q}$)



$13, 7, 1, 1, 1, 1$ ($\mathbb{Q}$)



$13, 6, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$



$13, 6, 2, 1, 1, 1$ $\left(\sqrt{-3}\right)$



$13, 5, 3, 1, 1, 1$ ($\mathbb{Q}$)



$13, 5, 2, 2, 1, 1\ \left(\sqrt{65}\right)$



$13, 5, 2, 2, 1, 1\ \left(\sqrt{65}\right)$



$13, 4, 3, 2, 1, 1\ (\mathbb{Q})$



$13, 3, 3, 2, 2, 1\ (\mathbb{Q})$



$12, 7, 2, 1, 1, 1\ \left(\sqrt{-3}\right)$



$12, 7, 2, 1, 1, 1\ \left(\sqrt{-3}\right)$



$12, 6, 3, 1, 1, 1\ (\mathbb{Q})$



$12, 6, 2, 2, 1, 1\ (\mathbb{Q})$



$12, 6, 2, 2, 1, 1\ (\mathbb{Q})$



$12, 5, 3, 2, 1, 1$ (quartic)



$12, 5, 3, 2, 1, 1$ (quartic)



$12, 5, 3, 2, 1, 1$ (quartic)



$12, 5, 3, 2, 1, 1$ (quartic)



$12, 5, 2, 2, 2, 1$ $(\mathbb{Q})$



$12, 4, 4, 2, 1, 1$ $(\mathbb{Q})$



$12, 4, 3, 3, 1, 1$ $(\mathbb{Q})$



$12, 4, 3, 3, 1, 1$ $(\mathbb{Q})$



$12, 4, 3, 2, 2, 1$ $(\mathbb{Q})$



$12, 4, 3, 2, 2, 1$ $\left(\sqrt{-3}\right)$



$12, 4, 3, 2, 2, 1$ $\left(\sqrt{-3}\right)$



$12, 3, 3, 3, 2, 1$ $(\mathbb{Q})$



$12, 3, 3, 2, 2, 2$ $(\mathbb{Q})$



$11, 9, 1, 1, 1, 1$ $(\mathbb{Q})$



$11, 8, 2, 1, 1, 1$ (cubic)



11, 8, 2, 1, 1, 1 (cubic)



11, 8, 2, 1, 1, 1 (cubic)



11, 7, 3, 1, 1, 1 (cubic)



11, 7, 3, 1, 1, 1 (cubic)



11, 7, 3, 1, 1, 1 (cubic)



11, 7, 2, 2, 1, 1 $\left(\sqrt{-7}\right)$



11, 7, 2, 2, 1, 1 $\left(\sqrt{-7}\right)$



11, 6, 4, 1, 1, 1 $\left(\sqrt{33}\right)$



11, 6, 4, 1, 1, 1 $\left(\sqrt{33}\right)$



11, 6, 3, 2, 1, 1 (cubic)



11, 6, 3, 2, 1, 1 (cubic)



11, 6, 3, 2, 1, 1 (cubic)



$11,5,5,1,1,1$ $(\mathbb{Q})$



$11,5,4,2,1,1$ (cubic)



$11,5,4,2,1,1$ (cubic)



$11,5,4,2,1,1$ (cubic)



$11,5,3,3,1,1$ $\left(\sqrt{5}\right)$



$11,5,3,3,1,1$ $\left(\sqrt{5}\right)$



$11,5,3,2,2,1$ $(\mathbb{Q})$



$11,4,4,3,1,1$ $(\mathbb{Q})$



$11,4,3,3,2,1$ $(\mathbb{Q})$



$10,10,1,1,1,1$ $(\mathbb{Q})$



$10,10,1,1,1,1$ $\left(\sqrt{5}\right)$



$10,10,1,1,1,1$ $\left(\sqrt{5}\right)$


$10, 9, 2, 1, 1, 1\ \left(\sqrt{5}\right)$


$10, 9, 2, 1, 1, 1\ \left(\sqrt{5}\right)$


$10, 8, 3, 1, 1, 1\ (\mathbb{Q})$


$10, 7, 4, 1, 1, 1\ (\mathbb{Q})$


$10, 7, 3, 2, 1, 1\ \left(\sqrt{21}\right)$


$10, 7, 3, 2, 1, 1\ \left(\sqrt{21}\right)$


$10, 7, 2, 2, 2, 1\ (\mathbb{Q})$


$10, 6, 5, 1, 1, 1\ (\mathbb{Q})$


$10, 6, 5, 1, 1, 1\ \text{(cubic)}$


$10, 6, 5, 1, 1, 1\ \text{(cubic)}$


$10, 6, 5, 1, 1, 1\ \text{(cubic)}$


$10, 6, 4, 2, 1, 1\ (\mathbb{Q})$



10, 6, 4, 2, 1, 1 (ℚ)



10, 6, 3, 2, 2, 1 (ℚ)



10, 5, 5, 2, 1, 1 $\left(\sqrt{5}\right)$



10, 5, 5, 2, 1, 1 $\left(\sqrt{5}\right)$



10, 5, 4, 2, 2, 1 (cubic)



10, 5, 4, 2, 2, 1 (cubic)



10, 5, 4, 2, 2, 1 (cubic)



10, 5, 3, 2, 2, 2 (ℚ)



10, 4, 4, 3, 2, 1 (ℚ)



10, 4, 3, 3, 2, 2 (ℚ)



9, 9, 2, 2, 1, 1 (ℚ)



9, 9, 2, 2, 1, 1 (cubic)



$9, 9, 2, 2, 1, 1$ (cubic)



$9, 9, 2, 2, 1, 1$ (cubic)



$9, 8, 3, 2, 1, 1$ (cubic)



$9, 8, 3, 2, 1, 1$ (cubic)



$9, 8, 3, 2, 1, 1$ (cubic)



$9, 7, 5, 1, 1, 1$ ($\mathbb{Q}$)



$9, 7, 4, 2, 1, 1$ $\left(\sqrt{-7}\right)$



$9, 7, 4, 2, 1, 1$ $\left(\sqrt{-7}\right)$



$9, 7, 3, 2, 2, 1$ (cubic)



$9, 7, 3, 2, 2, 1$ (cubic)



$9, 7, 3, 2, 2, 1$ (cubic)



$9, 6, 5, 2, 1, 1$ (cubic)



9, 6, 5, 2, 1, 1 (cubic)



9, 6, 5, 2, 1, 1 (cubic)



9, 6, 4, 3, 1, 1 ($\mathbb{Q}$)



9, 6, 3, 3, 2, 1 (cubic)



9, 6, 3, 3, 2, 1 (cubic)



9, 6, 3, 3, 2, 1 (cubic)



9, 5, 5, 2, 2, 1 ($\mathbb{Q}$)



9, 5, 4, 3, 2, 1 (cubic)



9, 5, 4, 3, 2, 1 (cubic)



9, 5, 4, 3, 2, 1 (cubic)



9, 5, 3, 3, 3, 1 ($\mathbb{Q}$)



9, 4, 3, 3, 3, 2 ($\mathbb{Q}$)



$8, 8, 4, 2, 1, 1$ ($\mathbb{Q}$)



$8, 8, 3, 3, 1, 1$ ($\mathbb{Q}$)



$8, 8, 3, 3, 1, 1$ $\left(\sqrt{-2}\right)$



$8, 8, 3, 3, 1, 1$ $\left(\sqrt{-2}\right)$



$8, 8, 2, 2, 2, 2$ ($\mathbb{Q}$)



$8, 7, 6, 1, 1, 1$ $\left(\sqrt{-3}\right)$



$8, 7, 6, 1, 1, 1$ $\left(\sqrt{-3}\right)$



$8, 7, 5, 2, 1, 1$ (cubic)



$8, 7, 5, 2, 1, 1$ (cubic)



$8, 7, 5, 2, 1, 1$ (cubic)



$8, 7, 4, 3, 1, 1$ $\left(\sqrt{-6}\right)$



$8, 7, 4, 3, 1, 1$ $\left(\sqrt{-6}\right)$



$8, 7, 4, 2, 2, 1\ \left(\sqrt{-7}\right)$



$8, 7, 4, 2, 2, 1\ \left(\sqrt{-7}\right)$



$8, 7, 3, 3, 2, 1\ (\mathbb{Q})$



$8, 6, 6, 2, 1, 1\ \left(\sqrt{-3}\right)$



$8, 6, 6, 2, 1, 1\ \left(\sqrt{-3}\right)$



$8, 6, 5, 3, 1, 1\ \left(\sqrt{5}\right)$



$8, 6, 5, 3, 1, 1\ \left(\sqrt{5}\right)$



$8, 6, 5, 2, 2, 1\ (\mathbb{Q})$



$8, 6, 4, 3, 2, 1\ \left(\sqrt{2}\right)$



$8, 6, 4, 3, 2, 1\ \left(\sqrt{2}\right)$



$8, 6, 4, 2, 2, 2\ (\mathbb{Q})$



$8, 5, 4, 3, 3, 1\ \left(\sqrt{10}\right)$



$8,5,4,3,3,1 \ \left(\sqrt{10}\right)$



$8,5,4,3,2,2 \ (\mathbb{Q})$



$8,4,4,4,3,1 \ (\mathbb{Q})$



$8,4,4,4,2,2 \ (\mathbb{Q})$



$7,7,7,1,1,1 \ (\mathbb{Q})$



$7,7,5,3,1,1 \ \left(\sqrt{21}\right)$



$7,7,5,3,1,1 \ \left(\sqrt{21}\right)$



$7,7,4,4,1,1 \ \left(\sqrt{-7}\right)$



$7,7,4,4,1,1 \ \left(\sqrt{-7}\right)$



$7,7,3,3,2,2 \ \left(\sqrt{7}\right)$



$7,7,3,3,2,2 \ \left(\sqrt{7}\right)$



$7,6,6,2,2,1 \ (\mathbb{Q})$



$7, 6, 5, 4, 1, 1$ $(\mathbb{Q})$



$7, 6, 5, 3, 2, 1$ (cubic)



$7, 6, 5, 3, 2, 1$ (cubic)



$7, 6, 5, 3, 2, 1$ (cubic)



$7, 6, 4, 4, 2, 1$ $(\mathbb{Q})$



$7, 5, 5, 4, 2, 1$ $\left(\sqrt{2}\right)$



$7, 5, 5, 4, 2, 1$ $\left(\sqrt{2}\right)$



$7, 5, 5, 3, 2, 2$ $(\mathbb{Q})$



$7, 5, 4, 4, 3, 1$ $(\mathbb{Q})$



$7, 5, 4, 3, 3, 2$ $(\mathbb{Q})$



$6, 6, 6, 4, 1, 1$ $(\mathbb{Q})$



$6, 6, 6, 2, 2, 2$ $(\mathbb{Q})$



$6,6,5,5,1,1\ \left(\sqrt{3}\right)$



$6,6,5,5,1,1\ \left(\sqrt{3}\right)$



$6,6,5,4,2,1\ (\mathbb{Q})$



$6,6,5,3,3,1\ (\mathbb{Q})$



$6,6,4,4,2,2\ (\mathbb{Q})$



$6,6,4,3,3,2\ (\mathbb{Q})$



$6,6,3,3,3,3\ (\mathbb{Q})$



$6,5,4,4,3,2\ (\mathbb{Q})$



$5,5,5,5,2,2\ (\mathbb{Q})$



$5,5,4,4,3,3\ (\mathbb{Q})$



$4,4,4,4,4,4\ (\mathbb{Q})$

Acknowledgements

We gratefully acknowledge many helpful comments from Alexander Hulpke, Wei Li and Will Sawin. YHH would like to thank the Science and Technology Facilities Council, UK, for an Advanced Fellowship and for STFC grant ST/J00037X/1, the Chinese Ministry of Education, for a Chang-Jiang Chair Professorship at NanKai University, the US NSF for grant CCF-1048082, as well as City University, London and Merton College, Oxford, for their enduring support. JM is grateful to the NSERC of Canada. JR is indebted to Oriel College and the Department of Physics, Oxford for support.

References

[1] John McKay and Abdellah Sebbar, "J-Invariants of Arithmetic Semistable Elliptic Surfaces and Graphs." *Proceedings on Moonshine and related topics (Montréal, QC, 1999)*, pages 119–130, 2001.

[2] Abdellah Sebbar, "Modular Subgroups, Forms, Curves and Surfaces." *Canad. Math. Bull.*, 45:294–308, 2002.

[3] Yang-Hui He and John McKay, "N=2 Gauge Theories: Congruence Subgroups, Coset Graphs and Modular Surfaces," arXiv:1201.3633 [hep-th].

[4] Jaap Top and Noriko Yui, "Explicit Equations of Some Elliptic Modular Surfaces." *Rocky Mountain J. Math.* Volume 37, Number 2 (2007), 663-687.

[5] Tetsuji Shioda, "On elliptic modular surfaces." *J. Math. Soc. Japan.* 24 (1972), 20-59.

[6] Alexander Zvonkin, "Functional Composition is a Generalized Symmetry." *Symmetry: Culture and Science*, 22:391–426, 2011.

[7] Leila Schneps (ed.), "The Grothendieck Theory of Dessins d'Enfants", LMS Lecture Note Series, CUP, ISBN 978-0-521-47821-2. 1994.

[8] Lando and Zvonkin, “Graphs on Surfaces and Their Applications”, Encyclopaedia of Mathematical Sciences: Lower-Dimensional Topology II, 141, Springer-Verlag, ISBN 978-3-540-00203-1, 2004.

[9] John Conway, Alexander Hulpke and John McKay, “On Transitive Permutation Groups.” *LMS J. Comput. Math 1*, pages 1–8, 1996.

[10] Rick Miranda and Ulf Persson, “Configurations of In Fibers on Elliptic K3-Surfaces.” *Math. Z.*, 201:339–361, 1989.

[11] Fritz Beukers and Hans Montanus, “Explicit calculation of elliptic fibrations of K3-surfaces and their Belyi-maps,” Number theory and polynomials, 33–51, LMS Lecture Note Series, 352, CUP, 2008.
Fritz Beukers, “Dessins Coming from the Miranda-Persson Table,” 2008, available at `http://www.staff.science.uu.nl/~beuke106/mirandapersson/Dessins.html`

[12] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.5.6*; 2012, `(\protect\vrule width0pt\protect\href{http://www.gap-system.org}{http://`

[13] The Sage Group, *Sage – Version 5.3*; 2012, `(\protect\vrule width0pt\protect\href{http`

[14] Derek Holt, Bettina Eick and Eamonn O’Brien. “Handbook of Computational Group Theory.” Discrete Mathematics and its Applications. Chapman and Hall, ISBN 978-1584883722, 2005.

[15] Marston Conder, “Schreier Coset Graphs and Their Applications.” *RIMS Kokyuroku.* 794, pages 169–175, 1992.

[16] David Morrison and Cumrun Vafa, “Compactifications of F Theory on Calabi-Yau Threefolds - I.” *Nucl. Phys. B*, 473, 1996.

[17] David Morrison and Cumrun Vafa, “Compactifications of F Theory on Calabi-Yau Threefolds - II.” *Nucl. Phys. B*, 476, 1996.

[18] Sujay Ashok, Freddy Cachazo and Eleonora Dell’Aquila, “Children’s Drawings From Seiberg-Witten Curves,” arXiv:1201.3633v2 [hep-th].

[19] Yang-Hui He, Wei Li, John McKay and James Read. *To appear.*